

# Complexity of First-Order Objective-Function-Free Optimization Algorithms Without a Bound on the Gradient

S. Gratton\*, S. Jerad† and Ph. L. Toint‡

4 I 2023

## Abstract

A class of algorithms for unconstrained nonconvex optimization is considered where the value of the objective function is never computed. The class contains a deterministic version of the first-order Adagrad method typically used for minimization of noisy function, but also allows the use of second-order information when available. The rate of convergence of methods in the class is analyzed and is shown to be identical to that known for first-order optimization methods using both function and gradients values. The result does not assume that the gradient of the objective function are uniformly bounded and is essentially sharp. It improves on previously known complexity bounds (in the stochastic context) by Défossez *et al.* (2020) and Gratton *et al.* (2022), both giving a better rate of global convergence and removing the need to assume bounded gradients. A new class of methods is designed, for which a slightly worse and essentially sharp complexity result holds. Limited numerical experiments show that the new methods' performance may be comparable to that of standard steepest descent, despite using significantly less information, and that this performance is relatively insensitive to noise.

**Keywords:** First-order methods, objective-function-free optimization (OFFO), Adagrad, convergence bounds, evaluation complexity, second-order models.

## 1 Introduction

This paper is concerned with Objective-Function-Free Optimization (OFFO<sup>(1)</sup>) algorithms, which we define as numerical optimization methods in which *the value of the problem's objective function is never calculated*, although we obviously assume that it exists. This is clearly at variance with the large majority of available numerical optimization algorithms, where the objective function is typically evaluated at every iteration, and its value then used to assess progress towards a minimizer and (often) to enforce descent. Dispensing with this information is therefore challenging. As it turns out, first-order OFFO methods (i.e. OFFO methods

---

\*Université de Toulouse, INP, IRIT, Toulouse, France. Email: serge.gratton@enseeiht.fr. Work partially supported by 3IA Artificial and Natural Intelligence Toulouse Institute (ANITI), French "Investing for the Future - PIA3" program under the Grant agreement ANR-19-PI3A-0004"

†ANITI, Université de Toulouse, INP, IRIT, Toulouse, France. Email: sadok.jerad@enseeiht.fr

‡NAXYS, University of Namur, Namur, Belgium. Email: philippe.toint@unamur.be. Partially supported by ANITI.

<sup>(1)</sup>This term is coined in contrast to DFO, a well-known acronym for Derivative Free Optimization. OFFO can be viewed as the complement of DFO, since the latter only uses objective function values and no derivatives, while the former only uses derivatives and no objective function values.

only using gradients) already exist for some time and have proved popular and useful in fields such as machine learning or sparse optimization, thereby justifying our interest. Algorithms such as Adagrad [12], RMSprop [28], Adam [21] or AMSgrad [27] have been proposed and analyzed, recent contributions being [10] and [31] where improved convergence bounds for such methods are discussed. All these results assume bounded gradients. We refer the reader to the interesting paper [10] for a more extensive historical perspective on the development of convergence theory of these first-order OFFO algorithms.

We take here the point of view that, although methods such as Adagrad and Adam have been defined and used in the context of probabilistic inexact gradient evaluations (such as resulting from sampling in finite-sum problems), they may nevertheless be of interest in a deterministic context (i.e. when gradients are computed exactly). Indeed, in addition to being one of the very few existing approaches to OFFO, their deterministic evaluation complexity analysis provides a very useful template for that of their probabilistic counterparts. In particular, lower complexity bounds for the deterministic case immediately apply to the stochastic one. The purpose of the present contribution is thus to explore the deterministic context, the probabilistic approach (using linear models only) being covered in a companion paper [18].

Should one be interested in first-order deterministic objective-function-free optimization, another obvious alternative would be to minimize the norm of the gradient (in the least-squares sense or in some other norm) using a derivative-free algorithm. To the best of our knowledge, this approach has not been widely experimented, maybe because it has the drawback of not being biased towards minimizing the underlying objective function: standard DFO algorithms, such as trust-region or adaptive regularization methods using finite-difference approximations, make, in this context, no distinction between minimizers, maximizers and other first-order points of the original problem. More importantly, convergence of the DFO minimizer would have to occur to a global minimum for the original gradient to approximately vanish. The evaluation complexity of this approach is nevertheless appealing under these strong assumptions, because it was proved in [4] (see also [15] and [6, Section 13.1]) that reaching an  $\epsilon$ -approximate first-order point (of the gradient's norm) can be obtained in  $\mathcal{O}(\epsilon^{-2} + |\log(\epsilon)|)$  evaluations<sup>(2)</sup>. We will compare below this theoretical bound with the new OFFO complexity results obtained in this paper.

**Contributions:** In this paper, we

1. interpret the deterministic counterpart of Adagrad as a first-order trust-region method and use this interpretation to extend it to potentially use second-order information,
2. provide, for this extended method and without assuming gradient boundeness, an essentially sharp global<sup>(3)</sup> bound on the gradient's norm as a function of the iteration counter, which is identical to that known for first-order optimization methods using both function and gradients values,
3. use the developed framework to define a further class of first-order OFFO methods, for which an essentially sharp complexity result is also provided,

---

<sup>(2)</sup>As is standard for two real positive sequences  $\{a_k\}$  and  $\{b_k\}$ , we say that  $a_k = \mathcal{O}(b_k)$  if and only if  $\lim_{k \rightarrow \infty} (a_k/b_k)$  is finite.

<sup>(3)</sup>I.e., valid at every iteration.

4. present some numerical experiments indicating that OFFO methods may indeed be competitive with steepest descent in efficiency and reliability.

Note: When this paper was completed, the authors became aware of the paper [16], where an adaptive trust-region algorithm is proposed in which the objective function is not evaluated, with a global rate of convergence similar to that presented below. However, while this algorithm is also a member of the class we study here, it is motivated differently (no relation to Adagrad is mentioned in [16]) and involves very different theoretical arguments (in particular, the question of sharpness for the complexity bound is not considered).

The paper is organized as follows. Section 2 introduces the new “trust-region minded” class of algorithms ASTR1 and the global rate of convergence of a subclass containing the deterministic Adagrad method are studied in Section 3. Section 4 then introduces a new ASTR1 subclass and analyzes its global rate of convergence. Some numerical illustration is provided in Section 5. Conclusions are finally outlined in Section 6.

## 2 A class of first-order minimization methods

We consider the problem

$$\min_{x \in \mathbb{R}^n} f(x) \tag{2.1}$$

where  $f$  is a smooth function from  $\mathbb{R}^n$  to  $\mathbb{R}$ . In particular, we will assume in what follows that

**AS.1:** the objective function  $f(x)$  is continuously differentiable;

**AS.2:** its gradient  $g(x) \stackrel{\text{def}}{=} \nabla_x^1 f(x)$  is Lipschitz continuous with Lipschitz constant  $L \geq 0$ , that is

$$\|g(x) - g(y)\| \leq L\|x - y\|$$

for all  $x, y \in \mathbb{R}^n$ , where, unless otherwise specified,  $\|\cdot\| = \|\cdot\|_2$  is the Euclidean norm;

**AS.3:** there exists a constant  $f_{\text{low}}$  such that, for all  $x$ ,  $f(x) \geq f_{\text{low}}$ .

AS.1, AS.2 and AS.3 are standard for the complexity analysis of optimization methods seeking first-order critical points, AS.3 guaranteeing in particular that the problem is well-posed. We stress once more that we do not assume that the gradient are uniformly bounded, at variance with [29, 10, 31, 18].

The class of methods of interest here are iterative and generate a sequence of iterates  $\{x_k\}_{k \geq 0}$ . The move from an iterate to the next directly depends on the gradient at  $x_k$  and algorithm-dependent *scaling factors*  $\{w_k = w(x_0, \dots, x_k)\}$  whose main purpose is to control the move’s magnitude. In our analysis, we will assume that

**AS.4:** for each  $i \in \{1, \dots, n\}$  there exists a constant  $\varsigma_i \in (0, 1]$  such that,  $w_{i,k} \geq \varsigma_i$  for all  $k \geq 0$ .

Since scaling factors are designed to control the length of the step, they are strongly reminiscent of the standard mechanism of the much studied trust-region optimization methods (see [8] for an extensive coverage and [30] for a more recent survey). In trust-region algorithms, a model of the objective function at an iterate  $x_k$  is built, typically using a truncated Taylor

series, and a step  $s_k$  is chosen that minimizes this model with a *trust-region*, that is a region where the model is assumed to represent the true objective function sufficiently well. This region is a ball around the current iterate, whose radius is updated adaptively from iteration to iteration, based on the quality of the prediction of the objective function value at the trial point  $x_k + s_k$ . For methods using gradient only, the model is then chosen as the first two terms of the Taylor’s expansion of  $f$  at the iterate  $x_k$ . Although, we are interested here in methods where the objective function’s value is not evaluated, and therefore cannot be used to accept/reject iterates and update the trust-region radius, a similar mechanism may be designed, this time involving the scaling factors  $\{w_k\}$ . The resulting algorithm, which we call ASTR1 (for Adaptively Scaled Trust Region using 1st order information) is stated on the following page.

The algorithm description calls for some comments.

1. Observe that we allow the use of second-order information by effectively defining a quadratic model

$$g_k^T s + \frac{1}{2} s^T B_k s \quad (2.10)$$

where  $B_k$  can of course be chosen as the true second-derivative matrix of  $f$  at  $x_k$  (provided it remains bounded to satisfy (2.3)) or any approximation thereof. Choosing  $B_k = 0$  results in a purely first-order algorithm.

The condition (2.3) on the Hessian approximations is quite weak, and allows in particular for a variety of quasi-Newton approaches, limited-memory or otherwise. In a finite-sum context, sampling bounded Hessians is also possible.

2. Conditions (2.5)–(2.8) define a “generalized Cauchy point”, much in the spirit of standard trust-region methodology (see [8, Section 6.3] for instance), where the quadratic model (2.10) is minimized in (2.8) along a good first-order direction ( $s_k^L$ ) to obtain a “Cauchy step”  $s_k^Q$ . Any step  $s_k$  can then be accepted provided it remains in the trust region (see (2.4)) and enforces a decrease in the quadratic model which is at least a fraction  $\tau$  of that achieved at the Cauchy step (see (2.5)).
3. At variance with many existing trust-region algorithms, the radius  $\Delta_k$  of the trust-region (2.2) is not recurred adaptively from iteration to iteration depending on how well the quadratic model predicts function values, but is directly defined as a scaled version of the local gradient. This is not without similarities with the trust-region method proposed by [13], which corresponds to a scaling factor equal to  $\|g_k\|^{-1}$ .
4. As stated, the ASTR1 algorithm does not include a termination rule, but such a rule can easily be introduced by terminating the algorithm in Step 1 if  $\|g_k\| \leq \epsilon$ , where  $\epsilon > 0$  is a user-defined first-order accuracy threshold.
5. It may seem to the reader that we have introduced two algorithmic parameters typically not present in existing OFFO methods. As it turns out, this is standard practice for trust-region methods and it is widely acknowledged that the behaviour of the algorithm is relatively insensitive to the choice made. Typically value are

$$\tau = \frac{1}{10} \quad \text{and} \quad \kappa_B = 10^5,$$

**Algorithm 2.1: ASTR1**

**Step 0: Initialization.** A starting point  $x_0$  is given. Constants  $\kappa_B \geq 1$  and  $\tau \in (0, 1]$  are also given. Set  $k = 0$ .

**Step 1: Define the TR.** Compute  $g_k = g(x_k)$  and define

$$\Delta_{i,k} = \frac{|g_{i,k}|}{w_{i,k}} \quad (2.2)$$

where  $w_k = w(x_0, \dots, x_k)$ .

**Step 2: Hessian approximation.** Select a symmetric Hessian approximation  $B_k$  such that

$$\|B_k\| \leq \kappa_B. \quad (2.3)$$

**Step 3: GCP.** Compute a step  $s_k$  such that

$$|s_{i,k}| \leq \Delta_{i,k} \quad (i \in \{1, \dots, n\}), \quad (2.4)$$

and

$$g_k^T s_k + \frac{1}{2} s_k^T B_k s_k \leq \tau \left( g_k^T s_k^Q + \frac{1}{2} (s_k^Q)^T B_k s_k^Q \right), \quad (2.5)$$

where

$$s_{i,k}^L = -\text{sgn}(g_{i,k}) \Delta_{i,k}, \quad (2.6)$$

$$s_k^Q = \gamma_k s_k^L, \quad (2.7)$$

with

$$\gamma_k = \begin{cases} \min \left[ 1, \frac{|g_k^T s_k^L|}{(s_k^L)^T B_k s_k^L} \right] & \text{if } (s_k^L)^T B_k s_k^L > 0, \\ 1 & \text{otherwise.} \end{cases} \quad (2.8)$$

**Step 4: New iterate.** Define

$$x_{k+1} = x_k + s_k, \quad (2.9)$$

increment  $k$  by one and return to Step 1.

the last one being possibly adapted to reflect the problem scaling. Note that these values are constant throughout the execution of the algorithm. At variance,  $\gamma_k$  is the iteration dependent stepsize, a quantity present in every first-order minimization method. Observe that we do not impose restrictions of the stepsize (beyond being positive), thereby covering most standard choices.

The algorithm being defined, the first step of our analysis is to derive a fundamental property of objective-function decrease, valid for all choices of the scaling factors satisfying AS.4.

**Lemma 2.1** Suppose that AS.1, AS.2 and AS.4 hold. Then we have that, for all  $k \geq 0$ ,

$$f(x_{j+1}) \leq f(x_j) - \sum_{i=1}^n \frac{\tau \varsigma_{\min} g_{i,j}^2}{2\kappa_B w_{i,j}} + \frac{1}{2}(\kappa_B + L) \sum_{i=1}^n \frac{g_{i,j}^2}{w_{i,j}^2} \quad (2.11)$$

and

$$f(x_0) - f(x_{k+1}) \geq \sum_{j=0}^k \sum_{i=1}^n \frac{g_{i,j}^2}{2\kappa_B w_{i,j}} \left[ \tau \varsigma_{\min} - \frac{\kappa_{\text{BBL}}}{w_{i,j}} \right] \quad (2.12)$$

where  $\varsigma_{\min} \stackrel{\text{def}}{=} \min_{i \in \{1, \dots, n\}} \varsigma_i$  and  $\kappa_{\text{BBL}} \stackrel{\text{def}}{=} \kappa_B(\kappa_B + L)$ .

**Proof.** Using (2.6) and AS.4, we deduce that, for every  $j \geq 0$ ,

$$|g_j^T s_j^L| = \sum_{i=1}^n \frac{g_{i,j}^2}{w_{i,j}} = \sum_{i=1}^n \frac{w_{i,j} g_{i,j}^2}{w_{i,j}^2} \geq \sum_{i=1}^n \frac{\varsigma_i g_{i,j}^2}{w_{i,j}^2} \geq \varsigma_{\min} \|s_j^L\|^2. \quad (2.13)$$

Suppose first that  $(s_j^L)^T B_j s_j^L > 0$  and  $\gamma_j < 1$ . Then, in view of (2.7), (2.8), (2.13) and (2.3),

$$g_j^T s_j^Q + \frac{1}{2}(s_j^Q)^T B_j s_j^Q = \gamma_j g_j^T s_j^L + \frac{1}{2} \gamma_j^2 (s_j^L)^T B_j s_j^L = -\frac{(g_j^T s_j^L)^2}{2(s_j^L)^T B_j s_j^L} \leq -\frac{\varsigma_{\min} |g_j^T s_j^L|}{2\kappa_B}.$$

Combining this inequality with the first equality in (2.13) then gives that

$$g_j^T s_j^Q + \frac{1}{2}(s_j^Q)^T B_j s_j^Q \leq -\frac{\varsigma_{\min}}{2\kappa_B} \sum_{i=1}^n \frac{g_{i,j}^2}{w_{i,j}}. \quad (2.14)$$

Suppose now that  $(s_j^L)^T B_j s_j^L \leq 0$  or  $\gamma_j = 1$ . Then, using (2.7), (2.14) and (2.6),

$$g_j^T s_j^Q + \frac{1}{2}(s_j^Q)^T B_j s_j^Q = g_j^T s_j^L + \frac{1}{2}(s_j^L)^T B_j s_j^L \leq \frac{1}{2} g_j^T s_j^L < 0$$

and (2.14) then again follows from the bound  $\kappa_B \geq 1$ . Successively using AS.1–AS.2, (2.5),

(2.14), (2.3) and (2.2) then gives that, for  $j \geq 0$ ,

$$\begin{aligned}
 f(x_{j+1}) &\leq f(x_j) + g_j^T s_j + \frac{1}{2} s_j^T B_j s_j - \frac{1}{2} s_j^T B_j s_j + \frac{1}{2} L \|s_j\|^2 \\
 &\leq f(x_j) + \tau \left( g_j^T s_j^Q + \frac{1}{2} (s_j^Q)^T B_j s_j^Q \right) + \frac{1}{2} (\kappa_B + L) \|s_j\|^2 \\
 &\leq f(x_j) - \sum_{i=1}^n \frac{\tau \varsigma_{\min} g_{i,j}^2}{2\kappa_B w_{i,j}} + \frac{1}{2} (\kappa_B + L) \sum_{i=1}^n \Delta_{i,j}^2 \\
 &\leq f(x_j) - \sum_{i=1}^n \frac{\tau \varsigma_{\min} g_{i,j}^2}{2\kappa_B w_{i,j}} + \frac{1}{2} (\kappa_B + L) \sum_{i=1}^n \frac{g_{i,j}^2}{w_{i,j}^2}
 \end{aligned}$$

This is (2.11). Summing up this inequality for  $j \in \{0, \dots, k\}$  then yields (2.12).  $\square$

Armed with Lemma 2.1, we are now in position to specify particular choices of the scaling factors  $w_{i,k}$  and derive the convergence properties of the resulting variants of ASTR1.

### 3 An Adagrad-like variant of ASTR1 using second-order models

We first consider a choice of scaling factors directly derived from the definition of the Adagrad algorithm [12]. For given  $\varsigma \in (0, 1]$ ,  $\vartheta \in (0, 1]$  and  $\mu \in (0, 1)$ , define, for all  $i \in \{1, \dots, n\}$  and for all  $k \geq 0$ ,

$$w_{i,k} \in \left[ \sqrt{\vartheta} v_{i,k}, v_{i,k} \right] \quad \text{where} \quad v_{i,k} \stackrel{\text{def}}{=} \left( \varsigma + \sum_{\ell=0}^k g_{i,\ell}^2 \right)^\mu. \quad (3.1)$$

The Adagrad scaling factors are recovered by  $\mu = \frac{1}{2}$  and  $\vartheta = 1$ , and ASTR1 with (3.1) and  $B_k = 0$  is then the standard (deterministic) Adagrad method. Our formulation with  $\mu \in (0, 1)$  allows a parametric analysis of methods “in the neighbourhood” of Adagrad, using first-order information or approximate second-order one. The  $\vartheta$  parameter is introduced for flexibility, in particular allowing non-monotone scaling factors<sup>(4)</sup>.

Before stating the global rate of convergence of the variant of ASTR1 using (3.1), we first prove a lemma, partly inspired by [29, 10].

**Lemma 3.1** Let  $\{a_k\}_{k \geq 0}$  be a non-negative sequence,  $\alpha > 0$ ,  $\xi > 0$  and define, for each  $k \geq 0$ ,  $b_k = \sum_{j=0}^k a_j$ . Then if  $\alpha \neq 1$ ,

$$\sum_{j=0}^k \frac{a_j}{(\xi + b_j)^\alpha} \leq \frac{1}{(1 - \alpha)} \left( (\xi + b_k)^{1-\alpha} - \xi^{1-\alpha} \right). \quad (3.2)$$

Otherwise (i.e. if  $\alpha = 1$ ),

$$\sum_{j=0}^k \frac{a_j}{(\xi + b_j)} \leq \log \left( \frac{\xi + b_k}{\xi} \right). \quad (3.3)$$

<sup>(4)</sup>Typical values are  $\varsigma_i = \frac{1}{100}$  and  $\vartheta = \frac{1}{1000}$ .

**Proof.** Consider first the case where  $\alpha \neq 1$  and note that  $\frac{1}{(1-\alpha)}x^{1-\alpha}$  is then a non-decreasing and concave function on  $(0, +\infty)$ . Setting  $b_{-1} = 0$  and using these properties, we obtain that, for  $j \geq 0$ ,

$$\begin{aligned} \frac{a_j}{(\xi + b_j)^\alpha} &\leq \frac{1}{1-\alpha} \left( (\xi + b_j)^{1-\alpha} - (\xi + b_j - a_j)^{1-\alpha} \right) \\ &\leq \frac{1}{1-\alpha} \left( (\xi + b_j)^{1-\alpha} - (\xi + b_{j-1})^{1-\alpha} \right). \end{aligned}$$

We then obtain (3.2) by summing this inequality for  $j \in \{0, \dots, k\}$ .

Suppose now that  $\alpha = 1$ , we then use the concavity and non-decreasing character of the logarithm to derive that

$$\frac{a_j}{(\xi + b_j)^\alpha} = \frac{a_j}{(\xi + b_j)} \leq \log(\xi + b_j) - \log(\xi + b_j - a_j) \leq \log(\xi + b_j) - \log(\xi + b_{j-1}).$$

The inequality (3.3) then again follows by summing for  $j \in \{0, \dots, k\}$ .  $\square$

From (3.2), we also obtain that, for  $\alpha < 1$ ,

$$\sum_{j=0}^k \frac{a_j}{(\xi + b_j)^\alpha} \leq \frac{1}{(1-\alpha)} (\xi + b_k)^{1-\alpha} \quad (3.4)$$

while, for  $\alpha > 1$ ,

$$\sum_{j=0}^k \frac{a_j}{(\xi + b_j)^\alpha} \leq \frac{\xi^{1-\alpha}}{(\alpha - 1)}. \quad (3.5)$$

Note that both the numerator and the denominator of the right-hand side of (3.2) tend to zero when  $\alpha$  tends to one. Applying l'Hospital rule, we then see that this right-hand side tends to the right-hand side of (3.3) and the bounds on  $\sum_{j=0}^k a_j / (\xi + b_j)^\alpha$  are therefore continuous at  $\alpha = 1$ .

Lemma 3.1 is crucial in the proof of our main complexity result, which we now state.



**Theorem 3.2** Suppose that AS.1–AS.3 hold and that the ASTR1 algorithm is applied to problem (2.1) with its scaling given by (3.1). If we define

$$\Gamma_0 \stackrel{\text{def}}{=} f(x_0) - f_{\text{low}},$$

then,

(i) if  $0 < \mu < \frac{1}{2}$ ,

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \frac{\kappa_1}{k+1}, \quad (3.6)$$

with

$$\kappa_1 = \max \left\{ \varsigma, \left[ \frac{4n\kappa_{\text{BBL}}}{(1-2\mu)\tau\varsigma^\mu\vartheta^{\frac{3}{2}}} \right]^{\frac{1}{\mu}}, \left[ \frac{2^{2\mu}\vartheta(1-2\mu)\Gamma_0}{n(\kappa_{\text{B}} + L)} \right]^{\frac{1}{1-2\mu}} \right\}; \quad (3.7)$$

(ii) if  $\mu = \frac{1}{2}$ ,

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \frac{\kappa_2}{k+1}, \quad (3.8)$$

with

$$\kappa_2 = \max \left\{ \varsigma, \frac{1}{2} e^{\frac{2\Gamma_0\vartheta}{n(\kappa_{\text{B}} + L)}}, \frac{1}{2\varsigma} \left( \frac{8n\kappa_{\text{B}}(\kappa_{\text{B}} + L)}{\tau\vartheta^{\frac{3}{2}}} \right)^2 \left| W_{-1} \left( -\frac{\tau\varsigma\vartheta^{\frac{3}{2}}}{8n\kappa_{\text{B}}(\kappa_{\text{B}} + L)} \right) \right|^2 \right\}, \quad (3.9)$$

where  $W_{-1}$  is the second branch of the Lambert function [9];

(iii) if  $\frac{1}{2} < \mu < 1$ ,

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \frac{\kappa_3}{k+1} \quad (3.10)$$

with

$$\kappa_3 \stackrel{\text{def}}{=} \max \left\{ \varsigma, \left[ \frac{2^{1+\mu}\kappa_{\text{B}}}{\tau\varsigma^\mu\sqrt{\vartheta}} \left( \Gamma_0 + \frac{n(\kappa_{\text{B}} + L)\varsigma^{1-2\mu}}{2\vartheta(2\mu-1)} \right) \right]^{\frac{1}{1-\mu}} \right\} \quad (3.11)$$

**Proof.** We see from (3.1) that  $w_{i,k}$  verifies **AS.4**. We may thus use Lemma 2.1. Moreover, (3.1) also implies that

$$\varsigma^\mu\sqrt{\vartheta} \leq w_{i,j} \leq \left( \varsigma + \sum_{\ell=0}^j \|g_\ell\|^2 \right)^\mu \quad (3.12)$$

for all  $j \geq 0$  and all  $i \in \{1, \dots, n\}$ . We now deduce from (2.2) and (2.12) that, for  $k \geq 0$ ,

$$f(x_{k+1}) \leq f(x_0) - \sum_{j=0}^k \frac{\tau\varsigma^\mu\sqrt{\vartheta} \|g_j\|^2}{2\kappa_{\text{B}} \max_{i \in \{1, \dots, n\}} w_{i,k}} + \frac{1}{2}(\kappa_{\text{B}} + L) \sum_{i=1}^n \sum_{j=0}^k \Delta_{i,j}^2. \quad (3.13)$$

For each  $i \in \{1, \dots, n\}$ , we then apply Lemma 3.1 with  $a_\ell = g_{i,\ell}^2$ ,  $\xi = \varsigma$  and  $\alpha = 2\mu < 1$ ,

and obtain from (2.2) and (3.1) that,

$$\sum_{j=0}^k \Delta_{i,j}^2 \leq \frac{1}{\vartheta(1-2\mu)} \left[ \left( \varsigma + \sum_{\ell=0}^k g_{i,\ell}^2 \right)^{1-2\mu} - \varsigma^{1-2\mu} \right] \leq \frac{1}{\vartheta(1-2\mu)} \left( \varsigma + \sum_{\ell=0}^k g_{i,\ell}^2 \right)^{1-2\mu}. \quad (3.14)$$

Now

$$\sum_{i=1}^n \sum_{j=0}^k \Delta_{i,j}^2 \leq \sum_{i=1}^n \frac{1}{\vartheta(1-2\mu)} \left( \varsigma + \sum_{i=1}^n \sum_{\ell=0}^k g_{i,\ell}^2 \right)^{1-2\mu} \leq \frac{n}{\vartheta(1-2\mu)} \left( \varsigma + \sum_{\ell=0}^k \|g_\ell\|^2 \right)^{1-2\mu} \quad (3.15)$$

and thus substituting this bound in (3.13) and using AS.3 gives that

$$\sum_{j=0}^k \frac{\tau \varsigma^\mu \sqrt{\vartheta} \|g_j\|^2}{2\kappa_B \max_{i \in \{1, \dots, n\}} w_{i,k}} \leq \Gamma_0 + \frac{n(\kappa_B + L)}{2\vartheta} (1-2\mu) \left( \varsigma + \sum_{j=0}^k \|g_j\|^2 \right)^{1-2\mu}. \quad (3.16)$$

Suppose now that

$$\sum_{j=0}^k \|g_j\|^2 \geq \max \left\{ \varsigma, \left[ \frac{2^{2\mu} \vartheta (1-2\mu) \Gamma_0}{n(\kappa_B + L)} \right]^{\frac{1}{1-2\mu}} \right\}, \quad (3.17)$$

implying

$$\varsigma + \sum_{j=0}^k \|g_j\|^2 \leq 2 \sum_{j=0}^k \|g_j\|^2 \quad \text{and} \quad \Gamma_0 \leq \frac{n(\kappa_B + L)}{2\vartheta(1-2\mu)} \left( 2 \sum_{j=0}^k \|g_j\|^2 \right)^{1-2\mu}.$$

Then, using (3.16) and (3.12),

$$\frac{\tau \varsigma^\mu \sqrt{\vartheta}}{2^{1+\mu} \kappa_B \left[ \sum_{\ell=0}^k \|g_\ell\|^2 \right]^\mu} \sum_{j=0}^k \|g_j\|^2 \leq \frac{2^{1-2\mu} n(\kappa_B + L)}{\vartheta(1-2\mu)} \left( \sum_{j=0}^k \|g_j\|^2 \right)^{1-2\mu}.$$

Solving this inequality for  $\sum_{j=0}^k \|g_j\|^2$  gives that

$$\sum_{j=0}^k \|g_j\|^2 \leq \left[ \frac{4n \kappa_{\text{BBL}}}{(1-2\mu)\tau \varsigma^\mu \vartheta^{\frac{3}{2}}} \right]^{\frac{1}{\mu}}$$

and therefore

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \left[ \frac{4n \kappa_{\text{BBL}}}{(1-2\mu)\tau \varsigma^\mu \vartheta^{\frac{3}{2}}} \right]^{\frac{1}{\mu}} \cdot \frac{1}{k+1}. \quad (3.18)$$

Alternatively, if (3.17) fails, then

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 < \max \left\{ \varsigma, \left[ \frac{2^{2\mu} \vartheta (1-2\mu) \Gamma_0}{2^{1-2\mu} n(\kappa_B + L)} \right]^{\frac{1}{1-2\mu}} \right\} \cdot \frac{1}{k+1}. \quad (3.19)$$

Combining (3.18) and (3.19) gives (3.6).

Let us now consider the case where  $\mu = \frac{1}{2}$ . For each  $i \in \{1, \dots, n\}$ , we apply Lemma 3.1 with  $a_k = g_{i,k}^2$ ,  $\xi = \varsigma$  and  $\alpha = 2\mu = 1$  and obtain that,

$$\sum_{i=1}^n \sum_{j=0}^k \Delta_{i,j}^2 \leq \frac{1}{\vartheta} \sum_{i=1}^n \log \left( \frac{1}{\varsigma} \left( \varsigma + \sum_{\ell=0}^k g_{i,\ell}^2 \right) \right) \leq \frac{n}{\vartheta} \log \left( 1 + \frac{1}{\varsigma} \sum_{\ell=0}^k \|g_\ell\|^2 \right).$$

and substituting this bound in (3.13) then gives that

$$\sum_{j=0}^k \frac{\tau \sqrt{\varsigma \vartheta} \|g_j\|^2}{2\kappa_B \max_{i \in \{1, \dots, n\}} w_{i,k}} \leq \Gamma_0 + \frac{n(\kappa_B + L)}{2\vartheta} \log \left( 1 + \frac{1}{\varsigma} \sum_{j=0}^k \|g_j\|^2 \right).$$

Suppose now that

$$\sum_{j=0}^k \|g_j\|^2 \geq \max \left[ \varsigma, \frac{1}{2} e^{\frac{2\vartheta \Gamma_0}{n(\kappa_B + L)}} \right], \quad (3.20)$$

implying that

$$1 + \frac{1}{\varsigma} \sum_{j=0}^k \|g_j\|^2 \leq \frac{2}{\varsigma} \sum_{j=0}^k \|g_j\|^2 \quad \text{and} \quad \Gamma_0 \leq \frac{n(\kappa_B + L)}{2\vartheta} \log \left( \frac{2}{\varsigma} \sum_{j=0}^k \|g_j\|^2 \right).$$

Using (3.12) for  $\mu = \frac{1}{2}$ , we obtain then that

$$\frac{\tau \sqrt{\varsigma \vartheta}}{2\sqrt{2} \kappa_B \sqrt{\sum_{\ell=0}^k \|g_\ell\|^2}} \sum_{j=0}^k \|g_j\|^2 \leq \frac{n(\kappa_B + L)}{\vartheta} \log \left( \frac{2}{\varsigma} \sum_{j=0}^k \|g_j\|^2 \right),$$

that is

$$\frac{\tau \sqrt{2\varsigma \vartheta}^{\frac{3}{2}}}{4\kappa_B} \sqrt{\sum_{j=0}^k \|g_j\|^2} \leq 2n(\kappa_B + L) \log \left( \sqrt{\frac{2}{\varsigma} \sum_{j=0}^k \|g_j\|^2} \right). \quad (3.21)$$

Now define

$$\gamma_1 \stackrel{\text{def}}{=} \frac{\tau \sqrt{2\varsigma \vartheta}^{\frac{3}{2}}}{4\kappa_B}, \quad \gamma_2 \stackrel{\text{def}}{=} 2n(\kappa_B + L) \quad \text{and} \quad u \stackrel{\text{def}}{=} \sqrt{\frac{2}{\varsigma} \sum_{j=0}^k \|g_j\|^2} \quad (3.22)$$

and observe that that  $\gamma_2 > 3\gamma_1$  because  $\tau \sqrt{\varsigma \vartheta}^{\frac{3}{2}} \leq 1$  and  $\kappa_B \geq 1$ . The inequality (3.21) can then be rewritten as

$$\gamma_1 u \leq \gamma_2 \log(u). \quad (3.23)$$

Let us denote by  $\psi(u) \stackrel{\text{def}}{=} \gamma_1 u - \gamma_2 \log(u)$ . Since  $\gamma_2 > 3\gamma_1$ , the equation  $\psi(u) = 0$  admits two roots  $u_1 \leq u_2$  and (3.23) holds for  $u \in [u_1, u_2]$ . The definition of  $u_2$  then gives that

$$\log(u_2) - \frac{\gamma_1}{\gamma_2} u_2 = 0$$

which is

$$u_2 e^{-\frac{\gamma_1}{\gamma_2} u_2} = 1.$$

Setting  $z = -\frac{\gamma_1}{\gamma_2} u_2$ , we obtain that

$$z e^z = -\frac{\gamma_1}{\gamma_2}$$

Thus  $z = W_{-1}\left(-\frac{\gamma_1}{\gamma_2}\right) < 0$ , where  $W_{-1}$  is the second branch of the Lambert function defined over  $[-\frac{1}{e}, 0)$ . As  $-\frac{\gamma_1}{\gamma_2} \geq -\frac{1}{3}$ ,  $z$  is well defined and thus

$$u_2 = -\frac{\gamma_2}{\gamma_1} z = -\frac{\gamma_2}{\gamma_1} W_{-1}\left(-\frac{\gamma_1}{\gamma_2}\right) > 0.$$

As a consequence, we deduce from (3.23) and (3.22) that

$$\sum_{j=0}^k \|g_j\|^2 = \frac{\varsigma}{2} u_2^2 = \frac{1}{2\varsigma} \left( \frac{8n\kappa_B(\kappa_B + L)}{\tau\vartheta^{\frac{3}{2}}} \right)^2 \left| W_{-1}\left(-\frac{\tau\varsigma\vartheta^{\frac{3}{2}}}{8n\kappa_B(\kappa_B + L)}\right) \right|^2.$$

and

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \frac{1}{2\varsigma} \left( \frac{8n\kappa_B(\kappa_B + L)}{\tau\vartheta^{\frac{3}{2}}} \right)^2 \left| W_{-1}\left(-\frac{\tau\varsigma\vartheta^{\frac{3}{2}}}{8n\kappa_B(\kappa_B + L)}\right) \right|^2 \cdot \frac{1}{k+1}. \quad (3.24)$$

If (3.20) does not hold, we have that

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 < \max \left\{ \varsigma, \frac{1}{2} e^{\frac{2\Gamma_0\vartheta}{n(\kappa_B + L)}} \right\} \cdot \frac{1}{k+1}. \quad (3.25)$$

Combining (3.24) and (3.25) gives (3.8).

Finally, suppose that  $\frac{1}{2} < \mu < 1$ . Once more, we apply Lemma 3.1 for each  $i \in \{1, \dots, n\}$  with  $a_\ell = g_{i,\ell}^2$ ,  $\xi = \varsigma$  and  $\alpha = 2\mu > 1$  and obtain that

$$\sum_{j=1}^k \Delta_{k,j}^2 \leq \frac{1}{\vartheta(1-2\mu)} \left( \left( \varsigma + \sum_{\ell=0}^k g_{i,\ell}^2 \right)^{1-2\mu} - \varsigma^{1-2\mu} \right) \leq \frac{\varsigma^{1-2\mu}}{\vartheta(2\mu-1)}. \quad (3.26)$$

Substituting the bound (3.26) in (3.13) and using (3.12) and AS.3 gives that

$$\sum_{j=0}^k \frac{1}{(\varsigma + \sum_{j=0}^k \|g_j\|^2)^\mu} \frac{\tau\varsigma^\mu \sqrt{\vartheta} \|g_j\|^2}{2\kappa_B} \leq \Gamma_0 + \frac{n(\kappa_B + L)\varsigma^{1-2\mu}}{2\vartheta(2\mu-1)}.$$

If we now suppose that

$$\sum_{j=0}^k \|g_j\|^2 \geq \varsigma, \quad (3.27)$$

then

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \left[ \frac{2^{1+\mu}\kappa_B}{\tau\varsigma^\mu \sqrt{\vartheta}} \left( \Gamma_0 + \frac{n(\kappa_B + L)\varsigma^{1-2\mu}}{2\vartheta(2\mu-1)} \right) \right]^{\frac{1}{1-\mu}} \cdot \frac{1}{k+1}. \quad (3.28)$$

If (3.27) does not hold, we derive that

$$\text{average}_{j \in \{0, \dots, k\}} \|g_j\|^2 \leq \frac{\varsigma}{(k+1)}. \quad (3.29)$$

Thus, (3.28) and (3.29) finally imply (3.10).  $\square$

These result suggest additional remarks.

1. [A comparison of the 3 bounds ??? Any suggestion?](#)
2. That the bounds given are not continuous as a function  $\mu$  at  $\mu = \frac{1}{2}$  is a result of our bounding process within the proof of Theorem 3.2 (for instance in the last inequality of (3.14)). Continuous bounds have been proved (see [17]) if one is ready to assume that the objective functions' gradients remain uniformly bounded.
3. If the algorithm is terminated as soon as  $\|g_k\| \leq \epsilon$  (which is customary for deterministic algorithms searching for first-order points), it must stop at the latest at iteration

$$k = \kappa_\star^2 \epsilon^{-2}, \quad (3.30)$$

where  $\kappa_\star = \kappa_1$  for  $\mu \in (0, \frac{1}{2})$ ,  $\kappa_\star = \kappa_2$  for  $\mu = \frac{1}{2}$  and  $\kappa_\star = \kappa_3$  for  $\mu \in (\frac{1}{2}, 1)$ . It is truly remarkable that there exist first-order OFFO methods whose global complexity order is identical to that of standard first-order methods using function evaluations (see [25, 19, 3] or [6, Chapter 2]), despite the fact that the latter exploit significantly more information.

4. If  $B_k = 0$  for all  $k$  and  $\mu = \frac{1}{2}$ , (3.8) gives an upper complexity bound for the deterministic momentum-less Adagrad algorithm which is significantly better than that proposed (using bounded gradients) by [10, 18] and more recently by [31] for a very specific choice of the stepsize (learning rate).
5. It is possible to give a more explicit bound on  $\kappa_2$  by finding an upper bound on the value of the involved Lambert function. This can be obtained by using [7, Theorem 1] which states that, for  $x > 0$ ,

$$|W_{-1}(-e^{-x-1})| \leq 1 + \sqrt{2x} + x. \quad (3.31)$$

Remembering that, for  $\gamma_1$  and  $\gamma_2$  given by (3.22),  $\log\left(\frac{\gamma_2}{\gamma_1}\right) \geq \log(3) > 1$  and taking  $x = \log\left(\frac{\gamma_2}{\gamma_1}\right) - 1 > 0$  in (3.31) then gives that

$$\left|W_{-1}\left(-\frac{\gamma_1}{\gamma_2}\right)\right| \leq \log\left(\frac{\gamma_2}{\gamma_1}\right) + \sqrt{2\left(\log\left(\frac{\gamma_2}{\gamma_1}\right) - 1\right)}.$$

6. It is also possible to extend the definition of  $s_k^L$  in (2.6) by premultiplying it by a stepsize  $\alpha_k \in [\alpha_{\min}, 1]$  for some  $\alpha_{\min} \in (0, 1]$ . Our results again remain valid (with modified constants). Covering a deterministic momentum-less Adam would require extending the results to allow for (3.1) to be replaced by

$$w_{i,k} = \varsigma + \sum_{j=0}^k \beta_2^{k-j} g_{i,j}^2 \quad (i \in \{1, \dots, n\}) \quad (3.32)$$

for some  $\beta_2 < 1$ . This can be done by following the argument of Theorem 2 in [10]. However, as in this reference, the final bound on the squared gradient norms does not tend to zero when  $k$  grows<sup>(5)</sup>, illustrating the (known) lack of convergence of Adam. We therefore do not investigate this option in detail.

7. The  $\epsilon$ -order (3.30) also improves slightly on the  $\mathcal{O}(\epsilon^{-2} + |\log(\epsilon)|)$   $\epsilon$ -order obtained for the DFO-based approach mentioned in the introduction (which, we recall, requires very strong assumptions on global minimization).

Are the (good) upper bound given by this theorem sharp?

**Theorem 3.3** The bounds (3.6), (3.8) and (3.10) are essentially sharp in that, for each  $\mu \in (0, 1)$  and each  $\eta \in (0, 1]$ , there exists a univariate function  $f_{\mu,\eta}$  satisfying AS.1-AS.3 such that, when applied to minimize  $f_{\mu,\eta}$  from the origin, the ASTR1 algorithm with (3.1) and  $\vartheta = 1$  produces a sequence of gradient norms given by  $\|g_k\| = \frac{1}{k^{\frac{1}{2}+\eta}}$ .

**Proof.** Following ideas of [6, Theorem 2.2.3], we first construct a sequence of iterates  $\{x_k\}$  for which  $f_{\mu,\eta}(x_k) = f_k$  and  $\nabla_x^1 f_{\mu,\eta}(x_k) = g_k$  for associated sequences of function and gradient values  $\{f_k\}$  and  $\{g_k\}$ , and then apply Hermite interpolation to exhibit the function  $f_{\mu,\eta}$  itself. We start by defining

$$g_0 \stackrel{\text{def}}{=} -2, \quad g_k \stackrel{\text{def}}{=} -\frac{1}{k^{\frac{1}{2}+\eta}} \quad (k > 0), \quad (3.33)$$

$$s_0 \stackrel{\text{def}}{=} \frac{2}{(\varsigma + 4)^\mu}, \quad s_k \stackrel{\text{def}}{=} \frac{1}{k^{\frac{1}{2}+\eta}(\varsigma + \sum_{j=0}^k g_j^2)^\mu} \quad (k > 0) \quad (3.34)$$

yielding that

$$|g_0 s_0| = \frac{4}{(\varsigma + 4)^\mu}, \quad |g_k s_k| = \frac{1}{k^{1+2\eta}(\varsigma + \sum_{j=0}^k g_j^2)^\mu} \leq \frac{1}{k^{1+2\eta}} \quad (k > 0) \quad (3.35)$$

(remember that  $g_0^2 = 4$ ). We then define  $B_k \stackrel{\text{def}}{=} 0$  for all  $k \geq 0$ ,

$$x_0 = 0, \quad x_{k+1} = x_k + s_k \quad (k > 0) \quad (3.36)$$

and

$$f_0 = \frac{4}{(\varsigma + 4)^\mu} + \zeta(1 + 2\eta) \quad \text{and} \quad f_{k+1} = f_k + g_k s_k \quad (k \geq 0), \quad (3.37)$$

where  $\zeta(\cdot)$  is the Riemann zeta function. Observe that the sequence  $\{f_k\}$  is decreasing and that, for all  $k \geq 0$ ,

$$f_{k+1} = f_0 - \sum_{k=0}^k |g_k s_k| \geq f_0 - \frac{4}{(\varsigma + 4)^\mu} - \sum_{k=1}^k \frac{1}{k^{1+2\eta}} \geq f_0 - \frac{4}{(\varsigma + 4)^\mu} - \zeta(1 + 2\eta)$$

---

<sup>(5)</sup>A constant term in  $-\log(\beta_2)$  refuses to vanish.

where we used (3.37) and (3.35). Hence (3.37) implies that

$$f_k \in [0, f_0] \text{ for all } k \geq 0. \quad (3.38)$$

Also note that, using (3.37),

$$|f_{k+1} - f_k - g_k s_k| = 0, \quad (3.39)$$

while, using (3.34),

$$|g_0 - g_1| = 1 \leq \frac{1}{2}(\varsigma + 4)^\mu s_0.$$

Moreover, using the fact that  $1/x^{\frac{1}{2}+\eta}$  is a convex function of  $x$  over  $[1, +\infty)$ , and that from (3.34)  $s_k \geq \frac{1}{k^{\frac{1}{2}+\eta}(\varsigma+4+k)^\mu}$ , we derive that, for  $k > 0$ ,

$$\begin{aligned} |g_{k+1} - g_k| &= \left| \frac{1}{(k+1)^{\frac{1}{2}+\eta}} - \frac{1}{k^{\frac{1}{2}+\eta}} \right| \\ &\leq \left( \frac{1}{2} + \eta \right) \frac{1}{k^{\frac{3}{2}+\eta}} \\ &\leq \frac{3}{2} \frac{(\varsigma + 4 + k)^\mu}{k k^{\frac{1}{2}+\eta} (\varsigma + 4 + k)^\mu} \\ &\leq \frac{3}{2} \frac{(\varsigma + 4 + k)^\mu}{k} s_k \\ &\leq \frac{3}{2} (\varsigma + 5)^\mu s_k. \end{aligned}$$

These last bounds and (3.38) allow us to use standard Hermite interpolation on the data given by  $\{f_k\}$  and  $\{g_k\}$ : see, for instance, Theorem A.9.1 in [6] with  $p = 1$  and

$$\kappa_f = \max \left[ \frac{3}{2}(\varsigma + 5)^\mu, f_0, 2 \right] \quad (3.40)$$

(the second term in the max bounding  $|f_k|$  because of (3.38) and the third bounding  $|g_k|$  because of (3.33)). We then deduce that there exists a continuously differentiable function  $f_{\mu,\eta}$  from  $\mathbb{R}$  to  $\mathbb{R}$  with Lipschitz continuous gradient (i.e. satisfying AS.1 and AS.2) such that, for  $k \geq 0$ ,

$$f_{\mu,\eta}(x_k) = f_k \quad \text{and} \quad \nabla_x^1 f_{\mu,\eta}(x_k) = g_k.$$

Moreover, the range of  $f_{\mu,\eta}$  and  $\nabla_x^1 f_{\mu,\eta}$  are constant independent of  $\eta$ , hence guaranteeing AS.3 and AS.3. The definitions (3.33), (3.34), (3.36) and (3.37) imply that the sequences  $\{x_k\}$ ,  $\{f_k\}$  and  $\{g_k\}$  can be seen as generated by the ASTR1 algorithm (with  $B_k = 0$ ) applied to  $f_{\mu,\eta}$ , starting from  $x_0 = 0$  and the desired conclusion follows.  $\square$

The bounds (3.6), (3.8) and (3.10) are therefore *essentially sharp* (in the sense of [5]) for the ASTR1 algorithm with (3.1) and  $\vartheta = 1$ , which is to say that the lower complexity bound for the algorithm is arbitrarily close to its upper bound. Interestingly, the argument in the proof of the above theorem fails for  $\eta = 0$ , as this choice yields that

$$\sum_{j=0}^k g_j^T s_j \geq \sum_{j=0}^k \frac{1}{k(\varsigma + \log(k+1))^\mu}.$$

Since

$$\int_1^k \frac{dt}{t(\log(t+1))^\mu} > \int_1^k \frac{dt}{(t+1)(\log(t+1))^\mu} = \frac{(\log(k+1))^{1-\mu}}{1-\mu} - \frac{\log(2)^{1-\mu}}{1-\mu}$$

tends to infinity as  $k$  grows, this indicates (in view (3.37)) that AS.3 cannot hold. Also note that (4.14) implies that the gradients remain uniformly bounded.

Figure 1 shows the behaviour of  $f_{\mu,\eta}(x)$  for  $\mu = \frac{1}{2}$  and  $\eta = \varsigma = \frac{1}{100}$ , its gradient and Hessian. The top three panels show the interpolated function resulting from the first 100 iterations of the ASTR1 algorithm with (3.1), while the bottom three panels report using  $10^4$  iterations. (We have chosen to shift  $f_0$  to 100 in order to avoid large numbers on the vertical axis of the left panels.) One verifies that the gradient is continuous and converges to zero. Since the Hessian remains bounded where defined, this indicates that the gradient is Lipschitz continuous. Due to the slow convergence of the series  $\sum_j 1/j^{1+2/100}$ , illustrating the boundeness of  $f_0 - f_{k+1}$  would require many more iterations. One also notes that the gradient is not monotonically increasing, which implies that  $f_{\mu,\eta}(x)$  is nonconvex, although this is barely noticeable in the left panels of the figure. Note finally that the fact that the example is unidimensional is not restrictive, since it is always possible to make the value of its objective function and gradient independent of all dimensions but one.




Figure 1: The function  $f_{\mu,\eta}(x)$  (left), its gradient  $\nabla_x^1 f_{\mu,\eta}(x)$  (middle) and its Hessian  $\nabla_x^2 f_{\mu,\eta}(x)$  (right) plotted as a function of  $x$ , for the first 100 (top) and  $10^4$  (bottom) iterations of the ASTR1 algorithm with (3.1) ( $\mu = \frac{1}{2}$ ,  $\eta = \varsigma = \frac{1}{100}$ )



## 4 A further “divergent stepsizes” variation on this theme

We now use a different proof technique to design new variants of ASTR1 with a fast global  $k$ -order. This is achieved by modifying the definition of the scaling factors  $w_{i,k}$ , requiring them to satisfy a fairly general growth condition explicitly depending on  $k$ , the iteration index. More specifically, we will assume, in this section, that the scaling factors  $w_{i,k}$  are chosen such that, for some power parameter  $0 < \nu \leq \mu < 1$ , all  $i \in \{1, \dots, n\}$  and some constants  $\varsigma_i \in (0, 1]$ ,

$$\max[\varsigma_i, v_{i,k}] (k+1)^\nu \leq w_{i,k} \leq \max[\varsigma_i, v_{i,k}] (k+1)^\mu \quad (k \geq 0), \quad (4.1)$$

where, for each  $i$ , the  $v_{i,k}$  satisfy the properties that

$$v_{i,k+1} > v_{i,k} \quad \text{implies that} \quad v_{i,k+1} \leq |g_{i,k+1}| \quad (4.2)$$

and

$$v_{i,k} \geq |g_{i,k}|/h(k) \quad (4.3)$$

for some positive function  $h(k)$  only depending on  $k$ . The motivation for introducing these new variants is the remarkable numerical performance [18] of particular choices where

$$v_{i,k} = \max_{j \in \{0, \dots, k\}} |g_{i,j}| \quad \text{and} \quad v_{i,k} = \frac{1}{k+1} \sum_{j \in \{0, \dots, k\}} |g_{i,j}|$$

which both satisfy (4.2) and (4.3) (with  $h(k) = 1$  for the first and  $h(k) = k+1$  for the second). We further illustrate this in Section 5.

We start by proving a useful technical result.

**Lemma 4.1** Consider an arbitrary  $i \in \{1, \dots, n\}$  and suppose that there exists a  $j_\varsigma$  such that

$$\min \left[ \frac{g_{i,j}^2}{\varsigma_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \leq \varsigma_i \quad \text{for } j \geq j_\varsigma. \quad (4.4)$$

Then

$$\min \left[ \frac{g_{i,j}^2}{\varsigma_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \geq \frac{g_{i,j}^2}{2\varsigma_i} \quad \text{for } j \geq j_\varsigma. \quad (4.5)$$

**Proof.** Suppose that there exists a  $j > j_\varsigma$  such that  $v_{i,j} > 2\varsigma_i$ . Assume, without loss of generality that  $j$  is the smallest such index. Then  $v_{i,j} > v_{i,j-1}$  and (4.2) implies that  $|g_{i,j}| \geq v_{i,j} \geq 2\varsigma_i$ . As a consequence,

$$\min \left[ \frac{g_{i,j}^2}{\varsigma_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \geq \min[4\varsigma_i, 2\varsigma_i] > \varsigma_i,$$

which contradicts (4.4). Thus no such  $j$  can exist and  $v_{i,j} \leq 2\varsigma_i$  for all  $j > j_*$  and (4.5) follows.  $\square$

We are now in position to state our complexity result for the ASTR1 algorithm using weight defined by (4.1), (4.2) and (4.3).

**Theorem 4.2** Suppose that AS.1, AS.2 and AS.3 hold and that the ASTR1 algorithm is applied to problem (2.1), where the scaling factors  $w_{i,k}$  are chosen in accordance with (4.1), (4.2) and (4.3). Then, for any  $\theta \in (0, \tau_{\zeta_{\min}})$  and

$$j_\theta \stackrel{\text{def}}{=} \left( \frac{\kappa_B(\kappa_B + L)}{\zeta_{\min}(\tau_{\zeta_{\min}} - \theta)} \right)^{\frac{1}{\nu}}, \quad (4.6)$$

there exist a constant  $\kappa_\diamond$ , a subsequence  $\{k_\ell\} \subseteq \{k_j\}_{j_\theta+1}^\infty$  and an index  $k_\zeta$  (where  $\kappa_\diamond$  and  $k_\zeta$  only depend on the problem and the algorithmic constants) such that, for all  $k_\ell \geq k_\zeta$ ,

$$\min_{j \in \{0, \dots, k_\ell\}} \|g_j\|^2 \leq \kappa_\diamond \frac{(k_\ell + 1)^\mu}{k_\ell - j_\theta} \leq \frac{2\kappa_\diamond(j_\theta + 1)}{k_\ell^{1-\mu}}. \quad (4.7)$$

**Proof.** From (2.12) and AS.3, using  $w_{\min,j} \stackrel{\text{def}}{=} \min_{i \in \{1, \dots, n\}} w_{i,k}$  ensures that

$$\Gamma_0 \geq f(x_0) - f(x_{k+1}) \geq \sum_{j=0}^k \sum_{i=1}^n \frac{g_{i,j}^2}{2\kappa_B w_{i,j}} \left[ \tau_{\zeta_{\min}} - \frac{\kappa_{\text{BBL}}}{w_{\min,j}} \right]. \quad (4.8)$$

Consider now an arbitrary  $\theta \in (0, \tau_{\zeta_{\min}})$  and suppose first that, for some  $j$ ,

$$\left[ \tau_{\zeta_{\min}} - \frac{\kappa_{\text{BBL}}}{w_{\min,j}} \right] \leq \theta, \quad (4.9)$$

i.e., using (4.1),

$$\zeta_{\min} j^\nu \leq w_{\min,j} \leq \frac{\kappa_{\text{BBL}}}{\tau_{\zeta_{\min}} - \theta}.$$

But this is impossible for  $j > j_\theta$  for  $j_\theta$  given by (4.6), and hence (4.9) fails for all  $j > j_\theta$ . As a consequence, we have that, for  $k > j_\theta$ ,

$$\begin{aligned} f(x_{j_\theta+1}) - f(x_k) &\geq \theta \sum_{j=j_\theta+1}^k \sum_{i=1}^n \frac{g_{i,j}^2}{2\kappa_B w_{i,j}} \\ &\geq \frac{\theta}{2\kappa_B} \sum_{j=j_\theta+1}^k \sum_{i=1}^n \frac{g_{i,j}^2}{\max[\zeta_i, v_{i,j}] (j+1)^\mu} \\ &\geq \frac{\theta}{2\kappa_B (k+1)^\mu} \sum_{j=j_\theta+1}^k \sum_{i=1}^n \min \left[ \frac{g_{i,j}^2}{\zeta_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \\ &\geq \frac{\theta(k-j_\theta)}{2\kappa_B (k+1)^\mu} \min_{j \in \{j_\theta+1, \dots, k\}} \left( \sum_{i=1}^n \min \left[ \frac{g_{i,j}^2}{\zeta_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \right) \end{aligned} \quad (4.10)$$

But we also know from (2.11), (4.1) and (4.3) that

$$\begin{aligned}
 f(x_0) - f(x_{j_\theta+1}) &\geq \sum_{j=0}^{j_\theta} \sum_{i=1}^n \frac{\tau_{\zeta_{\min}} g_{i,j}^2}{2\kappa_B w_{i,j}} - \frac{1}{2}(\kappa_B + L) \sum_{j=0}^{j_\theta} \sum_{i=1}^n \frac{g_{i,j}^2}{w_{i,j}^2} \\
 &\geq -\frac{1}{2}(\kappa_B + L) \sum_{j=0}^{j_\theta} \sum_{i=1}^n \frac{g_{i,j}^2}{w_{i,j}^2} \\
 &\geq -\frac{1}{2}(\kappa_B + L) \sum_{j=0}^{j_\theta} \sum_{i=1}^n \frac{g_{i,j}^2}{\max[\zeta, v_{i,k}]^2 (j+1)^{2\nu}} \\
 &\geq -\frac{1}{2}(\kappa_B + L) \sum_{j=0}^{j_\theta} \sum_{i=1}^n \frac{g_{i,j}^2}{v_{i,k}^2 (j+1)^{2\nu}} \\
 &\geq -\frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2. \tag{4.11}
 \end{aligned}$$

Combining (4.10) and (4.11), we obtain that

$$\Gamma_0 \geq f(x_0) - f(x_{k+1}) \geq -\frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 + \frac{\theta(k - j_\theta)}{2\kappa_B(k+1)^\mu} \min_{j \in \{j_\theta+1, \dots, k\}} \left( \sum_{i=1}^n \min \left[ \frac{g_{i,j}^2}{\zeta_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \right)$$

and thus that

$$\min_{j \in \{j_\theta+1, \dots, k\}} \left( \sum_{i=1}^n \min \left[ \frac{g_{i,j}^2}{\zeta_i}, \frac{g_{i,j}^2}{v_{i,j}} \right] \right) \leq \frac{2\kappa_B(k+1)^\mu}{\theta(k - j_\theta)} \left[ \Gamma_0 + \frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 \right]$$

and we deduce that there must exist a subsequence  $\{k_\ell\} \subseteq \{k\}_{j_\theta+1}^\infty$  such that, for each  $\ell$ ,

$$\sum_{i=1}^n \min \left[ \frac{g_{i,k_\ell}^2}{\zeta_i}, \frac{g_{i,jk_\ell}^2}{v_{i,k_\ell}} \right] \leq \frac{2\kappa_B(k_\ell+1)^\mu}{\theta(k_\ell - j_\theta)} \left[ \Gamma_0 + \frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 \right]. \tag{4.12}$$

But

$$\frac{(k_\ell+1)^\mu}{k_\ell - j_\theta} < \frac{2^\mu k_\ell^\mu}{k_\ell - j_\theta} < \frac{2k_\ell^\mu}{k_\ell - j_\theta} = \frac{2k_\ell^\mu k_\ell}{(k_\ell - j_\theta)k_\ell} = \frac{k_\ell}{k_\ell - j_\theta} \cdot \frac{2}{k_\ell^{1-\mu}} \leq \frac{2(j_\theta+1)}{k_\ell^{1-\mu}}, \tag{4.13}$$

where we used the facts that  $\mu < 1$  and that  $\frac{k_\ell}{k_\ell - j_\theta}$  is a decreasing function for  $k_\ell \geq j_\theta + 1$ . Using this inequality, we thus obtain from (4.12) that, for each  $\ell$ ,

$$\sum_{i=1}^n \min \left[ \frac{g_{i,k_\ell}^2}{\zeta_i}, \frac{g_{i,jk_\ell}^2}{v_{i,k_\ell}} \right] \leq \frac{4\kappa_B(j_\theta+1)}{\theta k_\ell^{1-\mu}} \left[ \Gamma_0 + \frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 \right].$$

As a consequence,

$$k_\zeta \stackrel{\text{def}}{=} \left( \frac{4\kappa_B(j_\theta+1) \left[ \Gamma_0 + \frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 \right]}{\theta \zeta_{\min}} \right)^{\frac{1}{1-\mu}}$$

is such that, for all  $k_\ell \geq k_\varsigma$ ,

$$\min \left[ \frac{g_{i,k_\ell}^2}{\varsigma_i}, \frac{g_{i,\ell}^2}{v_{i,k_\ell}} \right] \leq \varsigma_{\min}.$$

Lemma 4.1 then yields that, for all  $k_\ell \geq k_\varsigma$ ,

$$\sum_{i=1}^n \frac{g_{i,k_\ell}^2}{2\varsigma_i} \leq \frac{2\kappa_B(k_\ell + 1)^\mu}{\theta(k_\ell - j_\theta)} \left[ \Gamma_0 + \frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 \right]$$

which, because  $\varsigma_i \leq 1$ , gives that, for all  $k_\ell \geq k_\varsigma$ ,

$$\|g_{k_\ell}\|^2 \leq \frac{(k_\ell + 1)^\mu}{k_\ell - j_\theta} \left( \frac{4\kappa_B}{\theta} \right) \left[ \Gamma_0 + \frac{1}{2}n(\kappa_B + L) \sum_{j=0}^{j_\theta} h(j)^2 \right],$$

finally implying (4.7) because of (4.13).  $\square$

We again provide some comments on this last result.

1. The choice (4.1) is of course reminiscent, in a smooth and nonconvex setting, of the “divergent stepsize” subgradient method for non-smooth convex optimization (see [2] and the many references therein), for which a  $\mathcal{O}(1/\sqrt{k})$  global rate of convergence is known (Theorems 8.13 and 8.30 in this last reference).
2. Theorem 4.2 provides information on the speed of convergence for iterations that are beyond an *a priori* computable iteration index. Indeed  $j_\theta$  and  $k_\varsigma$  only depends on  $\nu$ ,  $h(\ell)$  and problem’s constants and, in particular do not depend on  $k$ . However, the formulation of the theorem is slightly weaker than that of Theorem 3.2. Because (4.7) only holds for iterates along the subsequence  $\{k_\ell\}$ , there is no guarantee that the bound given by the right-hand-side is valid at other iterations. But note that this right-hand side depends on  $k_\ell$ , which is an index in the complete sequence of iterates, rather than on  $\ell$  (the subsequence index).  
This slightly weaker formulation is no longer necessary if one is ready to assume bounded gradients, as can be seen in Theorem 4.1 in [17].
3. As the chosen values of  $\mu$  and  $\nu$  approach zero, then the  $k$ -order of convergence beyond  $j_\theta$  tends to  $\mathcal{O}(1/\sqrt{k_\ell})$ , which is the order derived for the methods of the previous section and is the standard  $k$ -order for first-order methods using evaluations of the objective function, albeit the value of  $j_\theta$  might increase.

We are now again interested to estimate how sharp the  $k$ -order bound (4.7) in  $\mathcal{O}(\frac{1}{k^{(1-\mu)/2}})$  is.

**Theorem 4.3** The bound (4.7) is essentially sharp in that, for any  $\omega > \frac{1}{2}(1 - \nu)$ , there exists a univariate function  $f_\omega(x)$  satisfying AS.1–AS.3 such that the ASTR1 algorithm with (4.1) applied to this function produces a sequence of gradient norms given by  $\|g_k\| = \frac{1}{(k+1)^\omega}$ .

**Proof.** Consider the sequence defined, for some  $\omega \in (\frac{1}{2}(1 - \nu), 1]$  and all  $k \geq 0$ , by

$$g_k = -\frac{1}{(k+1)^\omega} \quad w_k = \max \left[ \varsigma, \max_{\ell \in \{0, \dots, k\}} |g_\ell| \right] (k+1)^\nu = (k+1)^\nu, \quad (4.14)$$

$$s_k = \frac{1}{(k+1)^{2\omega-\nu}} < 1 \quad \text{and} \quad f_{k+1} = f_k + g_k s_k, \quad (4.15)$$

where we have chosen  $\varsigma \in (0, 1)$  and  $f_0 = \zeta(2\omega + \frac{1}{2})$  where  $\zeta(\cdot)$  is the Riemann zeta function. Immediately note that

$$\lim_{k \rightarrow \infty} |g_k| = 0,$$

and  $|g_k| \leq 1 = \kappa_g$  for all  $k$ . We now verify that, if

$$x_0 = 0 \quad \text{and} \quad x_k = x_{k-1} + s_{k-1} \quad \text{for } k \geq 1,$$

then exists a function  $f_\omega(x)$  satisfying AS.1–AS.3 such that, for all  $k \geq 0$ ,

$$f_\omega(x_k) = f_k, \quad \text{and} \quad g_\omega(x_k) = g_k,$$

and such that the sequence defined by (4.14)–(4.15) is generated by applying the ASTR1 algorithm using  $B_k = 0$ . The function  $f_\omega(x)$  is constructed using Hermite interpolation on each interval  $[x_k, x_{k+1}]$  (note that the  $x_k$  are monotonically increasing), which known (see [3] or [6, Th. A.9.2]) to exist whenever there exists a constant  $\kappa_f \geq 0$  such that, for each  $k$ ,

$$|f_{k+1} - f_k - g_k s_k| \leq \kappa_f |s_k|^2 \quad \text{and} \quad |g_{k+1} - g_k| \leq \kappa_f |s_k|.$$

The first of these conditions holds by construction of the  $\{f_k\}_{k \geq 0}$ . To verify the second, we first note that, because  $1/(k+1)^\omega$  is a convex function of  $k$  and  $|1/(k+1)| \leq 1$ ,

$$\frac{|g_{k+1} - g_k|}{|s_k|} \leq \frac{\omega(k+1)^{2\omega-\nu}}{(k+1)^{1+\omega}} \leq \frac{\omega}{(k+1)^{\nu-\omega+1}} \leq \omega \quad (k \geq 0), \quad (4.16)$$

where  $\nu - \omega + 1 \geq \nu > 0$ , so that the desired inequality holds with  $\kappa_f = \omega$ .

Moreover, Hermite interpolation guarantees that  $f_\omega(x)$  is bounded below whenever  $|f_k|$  and  $|s_k|$  remain bounded. We have already verified the second of these conditions in (4.15). We also have from (4.15) that

$$f_0 - f_{k+1} = \sum_{j=0}^k \frac{1}{(j+1)^{2\omega}(j+1)^\nu} \quad (4.17)$$

which converges to the finite limit  $\zeta(2\omega + \nu)$  because we have chosen  $\omega > \frac{1}{2}(1 - \nu)$ . Thus  $f_k \in (0, \zeta(2\omega + \nu)]$  for all  $k$  and the first condition is also satisfied and AS.3 holds. This completes our proof.  $\square$

The conclusions which can be drawn from this theorem parallel those drawn after Theorem 3.3. The bound (4.7) is essentially sharp (in the sense of [5]<sup>(6)</sup>) for the ASTR1 algorithm with (4.1).

<sup>(6)</sup>Observe that  $f_0$  now tends to infinity when  $\omega$  tends to  $\frac{1}{2}(\nu - 1)$  and hence that AS.3 fails in the limit. As before, the structure of (4.7) implies that the complexity bound deteriorates when the gap  $\Gamma_0 = f(x_0) - f_{\text{low}}$  grows.

Figure 2 shows the behaviour of  $f_\omega(x)$  for  $\nu = \frac{1}{9}$  and  $\omega = \frac{4}{9} + \frac{1}{100}$ , its gradient and Hessian. The top three panels show the interpolated function resulting from the first 100 iterations of the ASTR1 algorithm with (4.1), while the bottom three panels report using  $5 \cdot 10^4$  iterations. (We have again chosen to shift  $f_0$  to 100 in order to avoid large numbers on the vertical axis of the left panels.) As above, one verifies that the gradient is continous, non-monotone and converges to zero and that the Hessian remains bounded where defined, illustrating the gradient's Lipschitz continuity. Finally, as for Theorem 3.3, the argument in the proof of Theorem 4.3 fails for  $\omega = \frac{1}{2}(1 - \nu)$  because the sum in (4.17) is divergent in this case, which prevents AS.3 to hold.




Figure 2: The function  $f_\omega(x)$  (left), its gradient  $\nabla_x^1 f_\omega(x)$  (middle) and its Hessian  $\nabla_x^2 f_\omega(x)$  (right) plotted as a function of  $x$ , for the first 100 (top) and  $5 \cdot 10^4$  (bottom) iterations of the ASTR1 algorithm with (4.1) ( $\nu = \frac{1}{9}$ ,  $\omega = \frac{4}{9} + \frac{1}{100}$ )

## 5 Numerical illustration

Because the numerical behaviour of the methods discussed above is not well-known<sup>(7)</sup>, we now provide some numerical illustration. For the sake of clarity and conciseness, we needed to keep the list of algorithmic variants reported here reasonably limited, and have taken the following considerations into account for our choice.

1. Both scaling techniques (3.1) and (4.1) are illustrated. Moreover, since the Adam algorithm using (3.32) is so commonly used the stochastic context, we also included

<sup>(7)</sup>Except for stochastic finite sum minimization.

it in the comparison.

2. Despite Theorems 3.2 and 4.2 covering a wide choice of the parameters  $\mu$  and  $\nu$ , we have chosen to focus here on the most common choice for (3.1) and (3.32) (i.e.  $\mu = \frac{1}{2}$  and  $\beta_2 = \frac{9}{10}$ , corresponding to Adagrad and Adam). When using (4.1), we have also restricted our comparison to the single choice of  $\mu$  and  $\nu$  used (with reasonable success) in [18], namely  $\mu = \nu = \frac{1}{10}$ .
3. In order to be able to test enough algorithmic variants on enough problems in reasonable computing time, we have decided to limit our experiments to low-dimensional problems. For the same reason, we have focused our experiments on the case where  $\vartheta = 1$ .
4. We have chosen to define the step  $s_k$  in Step 3 of the ASTR1 algorithm by approximately minimizing the quadratic model (2.10) within the  $\ell_\infty$  trust-region using a projected truncated conjugate-gradient approach [23, 24] which is terminated as soon as

$$\|g_k + B_k s_k\|_2 \leq \max \left[ 10^{-12}, 10^{-5} \|g_k\|_2 \right].$$

We also considered an alternative, namely that of minimizing the quadratic model in an Euclidean  $\ell_2$  trust region (with the same accuracy requirement) using a Generalized Lanczos Trust Region (GLTR) technique [14].

5. We thought it would be interesting to compare “purely first-order” variants (that is variants for which  $B_k = 0$  for all  $k$ ) with methods using some kind of Hessian approximation. Among many possibilities, we selected three types of approximations of interest. The first is the diagonal Barzilai-Borwein approximation [1]

$$B_{k+1} = \frac{\|s_k\|_2^2}{y_k^T s_k} I_n \tag{5.1}$$

where  $I_n$  is the identity matrix of dimension  $n$ ,  $y_k = g_{k+1} - g_k$  and  $y_k^T s_k \geq 10^{-15} \|s_k\|_2^2$ . The second is limited-memory BFGS approximations [22], where a small number of BFGS updates are added to the matrix (5.1), each update corresponding to a secant pair  $(y_k, s_k)$ . The third is not to approximate the Hessian at all, but to use its exact value, that is  $B_k = \nabla_x^2 f(x_k)$  for all  $k$ .

Given these considerations, we have selected the following algorithmic variants:

**adagi1** : the ASTR1 algorithm using the Euclidean  $\|\cdot\|_2$  norm and

$$w_{i,k} = \left[ \varsigma + \sum_{j=0}^k \|g_j\|_2^2 \right]^{\frac{1}{2}} \quad \text{for } i \in \{1, \dots, n\},$$

**adagi1** : the ASTR1 algorithm using the  $\|\cdot\|_\infty$  norm and (3.1) with  $\mu = \frac{1}{2}$ ,

**adag2** : the ASTR1 algorithm using the Euclidean  $\|\cdot\|_2$  norm and

$$w_{i,k} = \left[ \varsigma + \sum_{j=0}^k \beta_2^{k-j} \|g_j\|_2^2 \right]^{\frac{1}{2}} \quad \text{for } i \in \{1, \dots, n\},$$

with  $\beta_2 = \frac{9}{10}$ ,

**adagi2** : the ASTR1 algorithm using the  $\|\cdot\|_\infty$  norm and (3.32) with  $\mu = \frac{1}{2}$  and

$\beta_2 = \frac{9}{10}$ ,

**maxg01** : the ASTR1 algorithm using (4.1) with

$$w_{i,k} = (k+1)^{\frac{1}{10}} \max \left[ \varsigma, \max_{j \in \{0, \dots, k\}} \|g_j\|_2 \right] \quad \text{for } i \in \{1, \dots, n\},$$

**maxgi01** : the ASTR1 algorithm using (4.1) with

$$w_{i,k} = (k+1)^{\frac{1}{10}} \max \left[ \varsigma, \max_{j \in \{0, \dots, k\}} |g_{i,j}| \right] \quad \text{for } i \in \{1, \dots, n\},$$

**sdba** : the standard steepest-descent algorithm using Armijo backtracking (see [6, Algorithm 2.2.1], for instance),

**b1adagi1** : **adagi1** where  $B_k$  is the Barzilai-Borwein Hessian approximation (5.1),

**lmadagi3b** : **adagi1** where  $B_k$  is a limited-memory BFGS Hessian approximation with 3 secant pairs,

**Eadagi1** : **adagi1** using the exact Hessian, i.e.  $B_k = \nabla^2 f(x_k)$  for all  $k$ .

When relevant, all variants use  $\varsigma = 0.01$ . The first seven algorithms are “purely first-order” in the sense discussed above. Note that, under AS.3, **maxg01** and **maxgi01** satisfy (4.1) with  $\mu = \nu = \frac{1}{10}$ ,  $\varsigma_i = \varsigma$  and  $\kappa_w = \kappa_g$ . Also note that **adagi1** and **adagi2** are nothing but the deterministic versions of Adagrad and Adam, respectively. All on the algorithms were run<sup>(8)</sup> on the low dimensional instances of the problems<sup>(9)</sup> of the OPM collection (January 2022) [20] listed with their dimension in Table 1, until either  $\|\nabla_x^1 f(x_k)\|_2 \leq 10^{-6}$ , or a maximum of 100000 iterations was reached, or evaluation of the gradient returned an error.

Before considering the results, we make two additional comments. The first is that very few of the test functions satisfy AS.3 on the whole of  $\mathbb{R}^n$ . While this is usually not a problem when testing standard first-order descent methods (because AS.3 may then be true in the level set determined by the starting point), this is no longer the case for significantly non-monotone methods like the ones tested here. As a consequence, it may (and does) happen that the gradient evaluation is attempted at a point where its value exceeds the Matlab overflow limit, causing the algorithm to fail on the problem. The second comment is that the

<sup>(8)</sup>In Matlab<sup>®</sup> on a Lenovo ThinkPad X1 Carbon with four cores and 8 GB of memory.

<sup>(9)</sup>From their standard starting point.



Problem	$n$	Problem	$n$	Problem	$n$	Problem	$n$	Problem	$n$	Problem	$n$
argauss	3	chebyqad	10	dixmaanl	12	heart8ls	8	msqrtals	16	scosine	10
arglina	10	cliff	2	dixon	10	helix	3	msqrtbls	16	sisser	2
arglinb	10	clplatea	16	dqartic	10	hilbert	10	morebv	12	spmsqrt	10
arglinc	10	clplateb	16	edensch	10	himln3	2	nlminsurf	16	tcontact	49
argtrig	10	clustr	2	eg2	10	himm25	2	nondquar	10	trigger	7
arwhead	10	cosine	10	eg2s	10	himm27	2	nzf1	13	tridia	10
bard	3	crglvy	4	eigfenals	12	himm28	2	osbornea	5	tlminsurf	16
bdarwhd	10	cube	2	eigenbls	12	himm29	2	osborneb	11	tnlminsurf	16
beale	2	curly10	10	eigencls	12	himm30	3	penalty1	10	vardim	10
biggs5	5	dixmaana	12	engvall	10	himm32	4	penalty2	10	vibrbeam	8
biggs6	6	dixmaanb	12	engval2	3	himm33	2	penalty3	10	watson	12
brownden	4	dixmaanc	12	expfit	2	hypcir	2	powellbs	2	wmsqrtals	16
booth	2	dixmaand	12	extrosnb	10	indef	10	powellsg	12	wmsqrtbls	16
box3	3	dixmaane	12	fminsurf	16	integreq	10	powellsq	2	woods	12
brkmcc	2	dixmaanf	12	freuroth	4	jensmp	2	powr	10	yfitu	3
brownal	10	dixmaang	12	genhumps	5	kowosb	4	recipe	2	zangwill2	2
brownbs	2	dixmaanb	12	gottfr	2	lminsurf	16	rosenbr	10	zangwill3	3
broyden3d	10	dixmaani	12	gulf	4	macino	10	sensors	10		
broydenbd	10	dixmaanj	12	hairy	2	mexhat	2	schmvett	3		
chandheu	10	dixmaank	12	heart6ls	6	meyer3	3	scurly10	10		

Table 1: The OPM test problems and their dimension

(sometimes quite wild) non-monotonicity of the methods considered here has another practical consequence: it happens on several nonconvex problems<sup>(10)</sup> that convergence of different algorithmic variants occurs to points with gradient norm within termination tolerance (the methods are thus achieving their objective), but these points can be quite far apart and have very different function values. It is therefore impossible to meaningfully compare the convergence performance to such points across algorithmic variants. This does reduce the set of problems where several variants can be compared.

We discuss the results of our tests from the efficiency and reliability points of view. Efficiency is measured in number of derivatives' evaluations (or, equivalently, iterations)<sup>(11)</sup>: the fewer evaluations the more efficient the algorithm. In addition to presenting the now standard performance profile [11] for our selection of algorithms in Figure 3, we follow [26] and consider the derived "global" measure  $\pi_{\text{algo}}$  to be  $\frac{1}{50}$  of the area below the curve corresponding to `algo` in the performance profile, for abscissas in the interval  $[1, 50]$ . The larger this area and closer  $\pi_{\text{algo}}$  to one, the closer the curve to the right and top borders of the plot and the better the global performance. When reporting reliability, we say that the run of an algorithmic variant on a specific test problem is successful if the gradient norm tolerance has been achieved, or if the final relative error on the objective-function value is below  $10^{-7}$  or, should the optimal value be below  $10^{-7}$ , if the final absolute error is below  $10^{-7}$ . These last two criteria were applied to instances of a total of 21 problems<sup>(12)</sup>. In what follows,  $\rho_{\text{algo}}$  denotes the percentage of successful runs taken on all problems where comparison is meaningful. Table 2 presents the values of these statistics in two columns: for easier reading, the variants are sorted by

<sup>(10)</sup>broyden3d, broydenbd, curly10, gottfr, hairy, indef, jensmp, osborneb, sensors, wmsqrtals, wmsqrtbls, woods.

<sup>(11)</sup>For `sdba`, gradient and objective-function evaluations.

<sup>(12)</sup>biggs6, brownden, box3, chebyqad, crglvy, cube, dixmaanb, dixmaanb, dixmaani, dixmaanb, dixmaanl, edensch, engval2, freuroth, indef, msqrtbls, osborneb, powellsq, rosenbr, vardim, zangwil3.

algo	$\pi_{\text{algo}}$	$\rho_{\text{algo}}$	algo	$\pi_{\text{algo}}$	$\rho_{\text{algo}}$
lmadagi3b	0.72	74.79	adagi1	0.70	75.63
adagi1	0.70	75.63	bladagi1	0.68	74.79
maxgi01	0.70	71.43	lmadagi3b	0.72	74.79
sdba	0.69	73.95	sdba	0.69	73.95
bladagi1	0.68	74.79	maxgi01	0.70	71.43
Eadagi1	0.60	67.23	adag1	0.54	68.91
adag1	0.54	68.91	Eadagi1	0.60	67.23
adag2	0.52	36.13	maxg01	0.52	63.03
maxg01	0.52	63.03	adag2	0.52	36.13
adagi2	0.51	31.09	adagi2	0.51	31.09

Table 2: Performance statistics for deterministic OFFO algorithms on OPM problems

decreasing global performance ( $\pi_{\text{algo}}$ ) in the first, and by decreasing reliability ( $\rho_{\text{algo}}$ ) in the second.




Figure 3: Performance profile for deterministic OFFO algorithms on OPM problems

A total of 22 problems<sup>(13)</sup> could not be successfully solved by any of the above algorithms, we believe mostly because of ill-conditioning.

The authors are of course aware that the very limited experiments presented here do not replace extended numerical practice and could be completed in various ways. They nevertheless suggest the following (very tentative) comments.

1. There seems to be a definite advantage in using the  $\|\cdot\|_{\infty}$  norm over  $\|\cdot\|_2$ , as can be seen by comparing `adag1` with `adagi1`, `adag2` with `adagi2`, and `maxg01` with `maxgi01`. While this may be due in part to the fact that the trust region in  $\ell_{\infty}$  norm is larger than that in  $\ell_2$  norm (and thus allows larger steps), it is also the case that the disaggregate

<sup>(13)</sup>biggs5, brownal, brownbs, cliff, eg2s, genhumps, gulf, heart8ls, himm29, mexhat, meyer3, nondquar, osbornea, penalty2, powellbs, powellsg, scurlly10, scosine, trigger, vibrbeam, watson, yfitu.

definition of the scaling factors  $w_{i,k}$  ((3.1), (3.32) or (4.1)) used in conjunction with the  $\ell_\infty$  norm may allow a better exploitation of differences of scale between coordinates.

2. Among the "purely first-order" methods, `sdba`, `maxgi01` and `adagi1` are almost undistinguishable from the performance point of view, with a small reliability advantage for `adagi1` (Adagrad). This means that, at least in those experiments, the suggestion resulting from the theory that OFFO methods may perform comparably to standard first-order methods seems vindicated.
3. The Adam variants (`adag2` and `adagi2`) are clearly outperformed in our tests by the Adagrad ones (`adag1` and `adagi1`). We recall that analytical examples where Adam fails do exist, while the convergence of Adagrad is covered by our theory.
4. The theoretical difference in global rate of convergence between `adagi1` and `maxgi01` does not seem to have much impact on the relative performance of these two methods.
5. The use of limited memory Hessian approximation (`lmadagi1`) appears to enhance the performance of `adagi1`, but this is not the base of the Barzilai-Borwein approximation (`bladagi1`) or, remarkably, for the use of the exact Hessian (`Eadagi1`). When these methods fail, this is often because the steplength is too small to allow the truncated conjugate-gradient solver to pick up second-order information in other directions than the negative gradient. What favours the limited memory approach remains unclear at this stage.

Finally, and although this is a slight digression from the paper's main topic, we report in Table 3 how reliability of our selection of OFFO variants is impacted by noise. To obtain these results, we ran the considered methods on all test problems where the evaluations (function<sup>(14)</sup> and derivatives) are contaminated by 5, 15, 25 or 50 % of relative Gaussian noise with unit variance. The reliability percentages in the table result from averaging ten sets of independent runs.

algo	$\rho_{\text{algo}}/\text{relative noise level}$				
	0%	5%	15%	25%	50%
<code>adagi1</code>	75.63	70.42	70.34	72.02	72.77
<code>bladagi1</code>	74.79	75.38	75.13	75.38	75.88
<code>lmadagi3b</code>	74.79	74.79	70.67	71.34	71.09
<code>sdba</code>	73.95	34.29	35.04	36.30	36.89
<code>maxgi01</code>	71.43	70.59	70.42	73.61	74.45
<code>adag1</code>	68.91	64.12	68.91	71.09	71.01
<code>Eadagi1</code>	67.23	67.98	68.74	70.17	70.08
<code>maxg01</code>	63.03	63.28	59.50	61.60	63.70
<code>adag2</code>	36.13	28.74	30.76	39.16	42.35
<code>adagi2</code>	31.09	24.12	25.88	28.66	31.93

Table 3: Reliability of OFFO algorithms as a function of the relative Gaussian noise level

<sup>(14)</sup>For `sdba`.

As can be seen in the table, the reliability of the `sdba` methods dramatically drops as soon as noise is present, while that of the other OFFO methods is barely affected and remains globally unchanged for increasing noise levels. This is consistent with widespread experience in the deep learning context, where noise is caused by sampling among the very large number of terms defining the objective function. This observation vindicates the popularity of methods such as Adagrad in the noisy context and suggests that the new OFFO algorithms may have some practical potential.

## 6 Conclusions

We have presented a parametric class of deterministic “trust-region minded” extensions of the Adagrad method, allowing for the use of second-order information, should it be available. We then prove that, for OFFO algorithms in this class,  $\min_{j \in \{0, \dots, k\}} \|g_j\| = \mathcal{O}(1/\sqrt{k+1})$ . We have shown that this bound, which does not require any uniform bound on the gradient, is essentially sharp. It is *identical to the global rate of convergence of standard first-order methods using both objective-function and gradient evaluations*, despite the fact that the latter exploit significantly more information. Thus, *if one considers the order of global convergence only, evaluating the objective-function values is an unnecessary effort*. We have also considered another class of OFFO algorithms inspired by the “divergent stepsize” paradigm in non-smooth convex optimization and have provided an essentially sharp (but slightly worse) global rate of convergence for this latter class. Limited numerical experiments suggest that the above theoretical conclusions may translate to practice and remain, for OFFO methods, relatively independent of noise.

Although discussed here in the context of unconstrained optimization, adaptation of the above OFFO algorithms to problems involving convex constraints (such as bounds on the variables) is relatively straightforward and practical: one then needs to intersect the trust-region with the feasible set and minimize the quadratic model in this intersection (see [8, Chapter 12]). It will be also of interest to further analyze the possible links between our proposals and those of [16], both from the theoretical and practical perspectives.

## References

- [1] J. Barzilai and J. Borwein. Two-point step size gradient method. *IMA Journal of Numerical Analysis*, 8:141–148, 1988.
- [2] A. Beck. *First-order Methods in Optimization*. Number 25 in MOS-SIAM Optimization Series. SIAM, Philadelphia, USA, 2017.
- [3] C. Cartis, N. I. M. Gould, and Ph. L. Toint. On the complexity of steepest descent, Newton’s and regularized Newton’s methods for nonconvex unconstrained optimization. *SIAM Journal on Optimization*, 20(6):2833–2852, 2010.
- [4] C. Cartis, N. I. M. Gould, and Ph. L. Toint. On the oracle complexity of first-order and derivative-free algorithms for smooth nonconvex minimization. *SIAM Journal on Optimization*, 22(1):66–86, 2012.
- [5] C. Cartis, N. I. M. Gould, and Ph. L. Toint. Worst-case evaluation complexity and optimality of second-order methods for nonconvex smooth optimization. In B. Sirakov, P. de Souza, and M. Viana, editors, *Invited Lectures, Proceedings of the 2018 International Conference of Mathematicians (ICM 2018), vol. 4, Rio de Janeiro*, pages 3729–3768. World Scientific Publishing Co Pte Ltd, 2018.
- [6] C. Cartis, N. I. M. Gould, and Ph. L. Toint. *Evaluation complexity of algorithms for nonconvex optimization*. Number 30 in MOS-SIAM Series on Optimization. SIAM, Philadelphia, USA, June 2022.

- [7] I. Chatzigeorgiou. Bounds on the Lambert function and their application to the outage analysis of user cooperation. *IEEE Communications Letters*, 17(8):1505–1508, 2013.
- [8] A. R. Conn, N. I. M. Gould, and Ph. L. Toint. *Trust-Region Methods*. Number 1 in MOS-SIAM Optimization Series. SIAM, Philadelphia, USA, 2000.
- [9] R. M. Corless, G. H. Gonnet, D. E. Hare, D. J. Jeffrey, and D. E. Knuth. On the Lambert W function. *Advances in Computational Mathematics*, 5:329–359, 1996.
- [10] A. Défossez, L. Bottou, F. Bach, and N. Usunier. A simple convergence proof for Adam and Adagrad. *Transactions on Machine Learning Research*, October 2022.
- [11] E. D. Dolan, J. J. Moré, and T. S. Munson. Optimality measures for performance profiles. *SIAM Journal on Optimization*, 16(3):891–909, 2006.
- [12] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12, July 2011.
- [13] J. Fan and Y. Yuan. A new trust region algorithm with trust region radius converging to zero. In D. Li, editor, *Proceedings of the 5th International Conference on Optimization: Techniques and Applications (ICOTA 2001, Hong Kong)*, pages 786–794, 2001.
- [14] N. I. M. Gould, S. Lucidi, M. Roma, and Ph. L. Toint. Solving the trust-region subproblem using the Lanczos method. *SIAM Journal on Optimization*, 9(2):504–525, 1999.
- [15] G. N. Grapiglia. Quadratic regularization methods with finite-difference gradient approximations. Optimization Online, November 2021.
- [16] G. N. Grapiglia and G. F. D. Stella. An adaptive trust-region method without function evaluation. *Computational Optimization and Applications*, 82:31–60, 2022.
- [17] S. Gratton, S. Jerad, and Ph. L. Toint. First-order objective-function-free optimization algorithms and their complexity. arXiv:2203.01757, 2022.
- [18] S. Gratton, S. Jerad, and Ph. L. Toint. Parametric complexity analysis for a class of first-order Adagrad-like algorithms. arXiv:2203.01647, 2022.
- [19] S. Gratton, A. Sartenaer, and Ph. L. Toint. Recursive trust-region methods for multiscale nonlinear optimization. *SIAM Journal on Optimization*, 19(1):414–444, 2008.
- [20] S. Gratton and Ph. L. Toint. OPM, a collection of optimization problems in Matlab. arXiv:2112.05636, 2021.
- [21] D. Kingma and J. Ba. Adam: A method for stochastic optimization. In *Proceedings in the International Conference on Learning Representations (ICLR)*, 2015.
- [22] D. C. Liu and J. Nocedal. On the limited memory BFGS method for large scale optimization. *Mathematical Programming, Series B*, 45(1):503–528, 1989.
- [23] J. J. Moré and G. Toraldo. Algorithms for bound constrained quadratic programming problems. *Numerische Mathematik*, 14:14–21, 1989.
- [24] J. J. Moré and G. Toraldo. On the solution of large quadratic programming problems with bound constraints. *SIAM Journal on Optimization*, 1(1):93–113, 1991.
- [25] Yu. Nesterov. *Introductory Lectures on Convex Optimization*. Applied Optimization. Kluwer Academic Publishers, Dordrecht, The Netherlands, 2004.
- [26] M. Porcelli and Ph. L. Toint. A note on using performance and data profiles for training algorithms. *ACM Transactions on Mathematical Software*, 45(2):1–25, 2019.
- [27] S. Reddi, S. Kale, and S. Kumar. On the convergence of Adam and beyond. In *Proceedings in the International Conference on Learning Representations (ICLR)*, 2018.
- [28] T. Tieleman and G. Hinton. Lecture 6.5-RMSPROP. COURSERA: Neural Networks for Machine Learning, 2012.
- [29] R. Ward, X. Wu, and L. Bottou. Adagrad stepsizes: sharp convergence over nonconvex landscapes. In *Proceedings in the International Conference on Machine Learning (ICML2019)*, 2019.
- [30] Y. Yuan. Recent advances in trust region algorithms. *Mathematical Programming, Series A*, 151(1):249–281, 2015.
- [31] D. Zhou, J. Chen, Y. Tang, Z. Yang, Y. Cao, and Q. Gu. On the convergence of adaptive gradient methods for nonconvex optimization. In *Proceedings of OPT2020: 12th Annual Workshop on Optimization for Machine Learning*, 2020.