

Complexity and performance for two classes of noise-tolerant first-order algorithms

S. Gratton*, S. Jerad† and Ph. L. Toint‡

20 IV 2022

Abstract

Two classes of algorithms for optimization in the presence of noise are presented, that do not require the evaluation of the objective function. The first generalizes the well-known Adagrad method. Its complexity is then analyzed as a function of its parameters, and it is shown that some methods of the class enjoy a better asymptotic convergence rate than previously known. A second class of algorithms is then derived whose complexity is at least as good as that of the first class. Initial numerical experiments on finite-sum problems arising from deep-learning applications suggest that that methods of the second class often outperform those of the first.

Keywords: First-order methods, objective-function-free optimization, noisy gradients, Adagrad, convergence bounds, evaluation complexity.

1 Introduction

Minimization algorithms which can handle noisy evaluations of the objective function and/or gradients have generated a significant amount of research in the last years [3, 5, 4, 6, 7, 8, 12, 13, 28, 14, 15, 21, 22, 25, 26, 27, 29, 31, 32]. Interestingly, a number of these contributions [3, 5, 4, 6, 7, 8, 12, 13, 25] indicate that, when the (noisy) objective function is evaluated, its accuracy is significantly more critical to ensure convergence than that of the computed (noisy) derivatives. This may be the reason why methods where the problem is avoided by *not* evaluating the objective function (such as Adagrad [15], RMSProp [27], Adam [21] or AMSGrad [26]) have become very popular in the context of finite-sum minimization, where noise in the evaluation arises from sampling among a very large number of terms. That such methods can be provably convergent to first-order stationary points is quite remarkable. Moreover, several authors have been able to prove global convergence rates, including the recent contributions by [14], where an improved (compared to earlier analysis) such rate was proved for the Adagrad algorithm, and by [31] where the analysis was refined to take

*Université de Toulouse, INP, IRIT, Toulouse, France. Email: serge.gratton@enseeiht.fr. Work partially supported by 3IA Artificial and Natural Intelligence Toulouse Institute (ANITI), French "Investing for the Future - PIA3" program under the Grant agreement ANR-19-PI3A-0004"

†ANITI, Université de Toulouse, INP, IRIT, Toulouse, France. Email: sadok.jerad@enseeiht.fr

‡NAXYS, University of Namur, Namur, Belgium. Email: philippe.toint@unamur.be. Partially supported by ANITI.

“sparsity” of the gradient sequence and optimal¹ learning rates into account and to cover AMSgrad and RMSProp.

The present paper uses (and extends) some preliminary results of [29, 14] to establish a new analysis that achieves several goals.

1. The global rate of convergence result of [14] is shown to hold for an extended class of methods comprising the Adagrad algorithm.
2. An improved asymptotic rate is also derived for these methods under an additional conditional variance condition akin to the Strong Growth Condition of [28], indicating that the results of [14] cannot be sharp if this condition holds. The new result is independent of gradient “sparsity”. More importantly, it allows an essentially arbitrary choice of the learning rate which *does not require the knowledge of the problem’s Lipschitz constant*. It therefore provides a strong alternative to those of [28] and [31].
3. Using the new analysis tools, a new class of methods is then proposed, whose global rate of convergence is shown to be very close to that of methods using (exact) function evaluations.
4. Numerical experiments with finite-sum problems arising from deep-learning applications indicate that method of the latter class often perform better than those of the former.

The presentation is organized as follows. A general framework of first-order trust-region algorithms is introduced in Section 2, in which two classes of algorithms (one of them containing the Adagrad method) are defined and analyzed (complexity-wise) in Sections 3 and 4, respectively. Numerical experiments in the finite-sum minimization context are presented in Section 5. Some conclusions are finally outlined in Section 6.

2 First-order trust-region methods for minimizing noisy functions

We are interested in (approximately) solving the problem

$$\min_{x \in \mathbb{R}^n} F(x) \tag{1}$$

where F is a function from \mathbb{R}^n to \mathbb{R} contaminated by noise. Moreover, we assume that evaluating F at any given x to sufficient accuracy is either impossible or too costly. Evaluating a noisy gradient is however possible. . . and our only source of information about the problem. While access to F or its exact gradient is impossible, we nevertheless make the following assumptions.

Assumption 2.1. The objective function $F(x)$ is continuously differentiable.

Assumption 2.2. Its exact gradient $G(x) \stackrel{\text{def}}{=} \nabla_x^1 f(x)$ is Lipschitz continuous with Lipschitz constant L , that is

$$\|G(x) - G(y)\| \leq L\|x - y\|$$

for all $x, y \in \mathbb{R}^n$.

¹A priori unknown.

Assumption 2.3. There exists a constant F_{low} such that, for all x , $F(x) \geq F_{\text{low}}$.

A standard consequence of Assumption 2.2 is that, for, any $x, s \in \mathbb{R}^n$,

$$F(x + s) \leq F(x) + G(x)^T s + \frac{L}{2} \|s\|^2 \quad (2)$$

(see Lemma 2.1 in [2] or Theorem A.8.3 in [11], for instance).

We now present a first-order *adaptively scaled gradient* algorithmic framework (ASGRAD), where, at iteration k , a *noisy* gradient $g_k = g(x_k)$ is evaluated and a step s_k defined that decreases the associated local linear model and whose size is determined by componentwise “scaling factors” $w_{i,k}$ to be chosen at each iteration. Our framework is formally described as follows.

Algorithm 2.1: The ASGRAD framework

Step 0: Initialization. x_0 and a constant $\gamma_{\text{low}} \in (0, 1]$ are given. Set $k = 0$.

Step 1: Step computation. Evaluate g_k and set

$$s_k = \gamma_k s_k^L, \quad (3)$$

with

$$s_{i,k}^L = -\frac{g_{i,k}}{w_{i,k}} \quad (4)$$

for a stepsize $\gamma_k \in [\gamma_{\text{low}}, 1]$ and positive scaling factors $w_{i,k}$.

Step 2: New iterate. Define

$$x_{k+1} = x_k + s_k, \quad (5)$$

increment k by one and return to Step 1.

We stress that g_k (as evaluated in Step 1) is a noisy random gradient evaluation. The algorithms of the ASGRAD framework therefore generate a stochastic process

$$\{x_k, g_k, \gamma_k, s_k^L, s_k\}$$

on some probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The associated expectation operator will be denoted by $\mathbb{E}[\cdot]$ and $\mathbb{E}_k[\cdot]$ will stand for the conditional expectation knowing $\{g_0, \dots, g_{k-1}\}$. All algorithms in our framework may clearly be interpreted as variants of Stochastic Gradient Descent, allowing for a variety of stepsize (learning rate) rules.

We will, in what follows, assume that the noisy gradient g_k is a bounded non-biased estimator of the true gradient, that is

Assumption 2.4. We have that, for all $k \geq 0$, $\mathbb{E}_k[g_k] = G(x_k)$. Moreover, there exists a constant $\kappa_g \geq 1$ such that $\|g_k\|_\infty \leq \kappa_g$ for all $k \geq 0$ and all realizations of the algorithm.

Note that this assumption, which is standard in the analysis of stochastic first-order methods, immediately implies that

$$\|G(x_k)\|_\infty \leq \kappa_g \quad (6)$$

for all $k \geq 0$.

The reader has undoubtedly noted that we have not been very specific regarding how the scaling factors $w_{i,k}$ are selected, and a whole range of options is possible. This justifies our choice to consider ASGRAD as an *algorithmic framework*, covering many possible such choices. The rest of this paper is devoted to the analysis of two specific classes of interest.

3 An Adagrad-inspired class of ASGRAD algorithms

In the first considered ASGRAD class, the scaling factors are inspired by the definition of the Adagrad algorithm [15]. More specifically, we make the following additional assumptions.

Assumption 3.1. For each $i \in \{1, \dots, n\}$ and $k \geq 0$, there exist a constant $\varsigma_i > 0$ and a random variable $v_{i,k}$ such that $v_{i,k} \geq \varsigma_i$ and $w_{i,k} = (v_{i,k})^\mu$ for some $\mu \in (0, 1)$. In addition,

$$|\mathbb{E}_k[v_{i,k}] - v_{i,k}| \leq \kappa_v (\mathbb{E}_k[g_{i,k}^2] + g_{i,k}^2) \quad (7)$$

for some $\kappa_v > 0$ and all $k \geq 0$.

Assumption 3.2. For every realization of the algorithm, we have that $g_{i,k}^2 \leq v_{i,k}$ for all $i \in \{1, \dots, n\}$ and all $k \geq 0$.

We immediately note that Assumption 3.1 implies that

$$v_{i,k} \geq \min_{i \in \{1, \dots, n\}} \varsigma_i \stackrel{\text{def}}{=} \varsigma_{\min} \quad (8)$$

and Assumption 3.2 ensures that

$$\mathbb{E}_k[g_{i,k}^2] \leq \mathbb{E}_k[v_{i,k}]. \quad (9)$$

The first step in our analysis is to derive a parametric bound on the decrease in the exact linear model of F caused by the step s_k , using a technique inspired by [29] and [14].

Lemma 3.3. Let s_j^L be the step produced at the j -th iteration by the an ASGRAD algorithm. Suppose also that Assumptions 2.4 3.1 and 3.2 hold. Let G_j be the true gradient of F at x_j . Then, for all $i \in \{1, \dots, n\}$,

$$\mathbb{E}_j[\gamma_j G_{i,j} s_{i,j}^L] \leq -\left(1 - \frac{\mu}{2}\right) \frac{\gamma_{\text{low}} G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + 2\kappa_\Delta \mathbb{E}_j\left[\frac{g_{i,j}^2}{w_{i,j}^2}\right], \quad (10)$$

where

$$\kappa_\Delta \stackrel{\text{def}}{=} \frac{\mu \kappa_v^2}{\gamma_{\text{low}}} \left[\kappa_g^{2\mu} + \frac{\kappa_g^2}{\varsigma_{\min}^{1-\mu}} + \frac{\kappa_g^{4-2\mu}}{\varsigma_{\min}^{2-2\mu}} + \kappa_g^{2-2\mu} \kappa_\mu \right] \quad \text{with} \quad \kappa_\mu \stackrel{\text{def}}{=} \frac{1}{\varsigma_{\min}^{1-2\mu}} \mathbf{1}_{\mu < \frac{1}{2}} + \kappa_g^{4\mu-2} \mathbf{1}_{\mu \geq \frac{1}{2}}, \quad (11)$$

where $\mathbf{1}_e$ stands for the indicator function of the event e .

Proof. See Appendix.

This lemma essentially implies that s^L provides a descent direction on the true F as long as the square of the true gradient's norm remains large compared with the stepsizes. We also need another result, partly inspired by [29, 14], whose utility will be to bound the last term on the right-hand side of (10).

Lemma 3.4. *Let $\{a_k\}_{k \geq 0}$ be a non-negative sequence, $\alpha > 0$ and define, for each $k \geq 0$, $b_k = \sum_{j=0}^k a_j$. Then if $\alpha \neq 1$,*

$$\sum_{j=0}^k \frac{a_j}{(\varsigma + b_j)^\alpha} \leq \frac{1}{(1-\alpha)} ((\varsigma + b_k)^{1-\alpha} - \varsigma^{1-\alpha}). \quad (12)$$

Otherwise (i.e. if $\alpha = 1$) (see Lemma 5.2 in [14]),

$$\sum_{j=0}^k \frac{a_j}{\varsigma + b_j} \leq \log \left(\frac{\varsigma + b_k}{\varsigma} \right). \quad (13)$$

Proof. See Appendix. Note that (13) is the limit of (12) when α tends to one.

Using both Lemmas 3.3 and 3.4, we are now in position to deduce a first result on the global convergence rate of a class of ASGRAD algorithms using specific ‘‘Adagrad-like’’ scaling factors satisfying Assumptions 3.1 and 3.2.

Theorem 3.5. *Suppose that Assumptions 2.1–2.4 hold and that an ASGRAD algorithm is applied to problem (1) where, for all $k \geq 0$ and all $i \in \{1, \dots, n\}$,*

$$w_{i,k} = \left(\varsigma + \sum_{\ell=0}^k g_{i,\ell}^2 \right)^\mu, \quad (14)$$

where $\varsigma \in (0, \kappa_g]$ and $\mu \in (0, 1)$. Then the following bounds hold for κ_Δ given in (11) and

$$\kappa_\square \stackrel{\text{def}}{=} \frac{\kappa_g^{2\mu} (4\kappa_\Delta + L)}{(1 - \frac{\mu}{2}) \gamma_{\text{low}}}. \quad (15)$$

(i) *If $\mu \in (0, \frac{1}{2})$, then*

$$\begin{aligned} \mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] &\leq \frac{2\kappa_g^{2\mu}}{(1 - \frac{\mu}{2}) \gamma_{\text{low}} (k+1)^{1-\mu}} [F(x_0) - F_{\text{low}}] \\ &\quad + \frac{n\kappa_\square}{1-2\mu} \frac{(\varsigma + \kappa_g^2(k+1))^{1-2\mu} - \varsigma^{1-2\mu}}{(k+1)^{1-\mu}}. \end{aligned} \quad (16)$$

(ii) *If $\mu = \frac{1}{2}$, then*

$$\mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq \frac{8\kappa_g}{3\gamma_{\text{low}} \sqrt{(k+1)}} [F(x_0) - F_{\text{low}}] + n\kappa_\square \frac{\log \left(1 + (k+1) \frac{\kappa_g^2}{\varsigma} \right)}{\sqrt{(k+1)}}. \quad (17)$$

(iii) If $\mu \in (\frac{1}{2}, 1)$, then

$$\begin{aligned} \mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] &\leq \frac{2\kappa_g^{2\mu}}{(1 - \frac{\mu}{2})\gamma_{\text{low}}(k+1)^{1-\mu}} \left[F(x_0) - F_{\text{low}} \right] \\ &\quad + \frac{n\kappa_{\square}}{2\mu - 1} \frac{\varsigma^{1-2\mu} - (\varsigma + \kappa_g^2(k+1))^{1-2\mu}}{(k+1)^{1-\mu}}. \end{aligned} \quad (18)$$

Proof. It is clear from (14) that $w_{i,k} \geq \varsigma^\mu$. Moreover, if we define $v_{i,k} \stackrel{\text{def}}{=} \varsigma + \sum_{\ell=0}^k g_{i,\ell}^2$, then we have that $w_{i,k} = v_{i,k}^\mu$, $v_{i,k} \geq g_{i,k}^2$ and

$$|\mathbb{E}_k[v_{i,k}] - v_{i,k}| = |\mathbb{E}_k[g_{i,k}^2] - g_{i,k}^2| \leq \mathbb{E}_k[g_{i,k}^2] + g_{i,k}^2.$$

Thus the proposed scaling factors verify Assumptions 3.1 and 3.2 with $\kappa_v = 1$.

Using (2), we derive that

$$F(x_{j+1}) \leq F(x_j) + \gamma_j G_j^T s_j^L + \frac{L}{2} \gamma_j^2 \|s_j^L\|^2 \leq F(x_j) + \gamma_j G_j^T s_j^L + \frac{L}{2} \|s_j^L\|^2.$$

Taking the conditional expectation, using Lemma 3.3, the fact that $v_{i,j} \leq (k+2)\kappa_g^2$ (because we assumed that $\varsigma \leq \kappa_g$), (4), we deduce that, for $j \in \{0, \dots, k\}$,

$$\begin{aligned} \mathbb{E}_j[F(x_{j+1})] &\leq F(x_j) + \sum_{i=1}^n \mathbb{E}_j[\gamma_j G_{i,j} s_{i,j}^L] + \frac{L}{2} \mathbb{E}_j[\|s_j^L\|^2], \\ &\leq F(x_j) - \sum_{i=1}^n \left(1 - \frac{\mu}{2}\right) \gamma_{\text{low}} \frac{G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + 2\kappa_\Delta \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j}^2} \right] + \frac{L}{2} \mathbb{E}_j[\|s_j^L\|^2], \\ &\leq F(x_j) - \left(1 - \frac{\mu}{2}\right) \gamma_{\text{low}} \frac{\|G_j\|^2}{\kappa_g^{2\mu}(k+2)^\mu} + \left(\frac{L}{2} + 2\kappa_\Delta\right) \mathbb{E}_j[\|s_j^L\|^2]. \end{aligned}$$

We may now sum the previous inequality for $j \in \{0, \dots, k\}$ and take the full expectation to derive that

$$\begin{aligned} \mathbb{E}[F(x_{k+1})] &\leq F(x_0) - \left(1 - \frac{\mu}{2}\right) \frac{\gamma_{\text{low}}}{\kappa_g^{2\mu}(k+2)^\mu} \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] + \left(\frac{L}{2} + 2\kappa_\Delta\right) \sum_{j=0}^k \mathbb{E}[\|s_j^L\|^2] \\ &\leq F(x_0) - \left(1 - \frac{\mu}{2}\right) \frac{\gamma_{\text{low}}}{\kappa_g^{2\mu}(k+2)^\mu} \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] + \left(\frac{L}{2} + 2\kappa_\Delta\right) \sum_{i=1}^n \sum_{j=0}^k \mathbb{E}[(s_{i,j}^L)^2]. \end{aligned} \quad (19)$$

Using now Lemma 3.4 with $\alpha = 2\mu$ for each $s_{i,j}^L$, (4), (14) and Assumption 2.4, we derive that, for $\mu \in (0, \frac{1}{2})$,

$$\begin{aligned} \sum_{j=0}^k (s_{i,j}^L)^2 &= \sum_{j=0}^k \frac{g_{i,j}^2}{(\varsigma + \sum_{j=0}^k g_{i,j}^2)^{2\mu}} \\ &\leq \frac{1}{1 - 2\mu} \left[\left(\varsigma + \sum_{j=0}^k g_{i,j}^2 \right)^{1-2\mu} - \varsigma^{1-2\mu} \right] \\ &\leq \frac{1}{1 - 2\mu} \left[\left(\varsigma + (k+1)\kappa_g^2 \right)^{1-2\mu} - \varsigma^{1-2\mu} \right]. \end{aligned}$$

Plugging this inequality in (19) and using Assumption 2.3, we obtain that

$$F_{\text{low}} \leq \mathbb{E}[F(x_{k+1})] \leq F(x_0) - (1 - \frac{\mu}{2}) \frac{\gamma_{\text{low}}}{\kappa_g^{2\mu} (k+2)^\mu} \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] \\ + \frac{n}{1-2\mu} \left(\frac{L}{2} + 2\kappa_\Delta \right) [(\varsigma + (k+1)\kappa_g^2)^{1-2\mu} - \varsigma^{1-2\mu}]$$

and thus, since $(k+2)^\mu \leq 2(k+1)^\mu$, that

$$(k+1) \mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] \quad (20)$$

$$\leq \frac{2\kappa_g^{2\mu} (F(x_0) - F_{\text{low}})}{(1 - \frac{\mu}{2}) \gamma_{\text{low}} (k+1)^{-\mu}} \quad (21) \\ + \frac{n [(\varsigma + \kappa_g^2(k+1))^{1-2\mu} - \varsigma^{1-2\mu}]}{(1-2\mu)(k+1)^{-\mu}} \left(\frac{\kappa_g^{2\mu} (L + 4\kappa_\Delta)}{\gamma_{\text{low}} (1 - \frac{\mu}{2})} \right),$$

which is (16).

If $\mu = \frac{1}{2}$, we reuse (19) and Lemma 3.4 for each $s_{i,j}^L$ with $\alpha = 1$, and derive that, in this case,

$$\mathbb{E}[F(x_{k+1})] \leq F(x_0) - \frac{3}{4} \frac{\gamma_{\text{low}}}{\sqrt{(k+2)\kappa_g}} \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] + n \left(\frac{L}{2} + 2\kappa_\Delta \right) \log \left(1 + (k+1) \frac{\kappa_g^2}{\varsigma} \right).$$

By a reasoning similar to that leading to (20) we now obtain that

$$(k+1) \mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] \\ \leq \left(\frac{4}{3} \right) \frac{2\kappa_g (F(x_0) - F_{\text{low}}) \sqrt{(k+1)}}{\gamma_{\text{low}}} \\ + \left(\frac{4n}{3} \right) \frac{\kappa_g}{\gamma_{\text{low}}} (L + 4\kappa_\Delta) \log \left(1 + (k+1) \frac{\kappa_g^2}{\varsigma} \right) \sqrt{(k+1)}.$$

Rearranging the terms yields (17).

Finally, if $\mu \in (\frac{1}{2}, 1)$, we again reuse (19) and Lemma 3.4 for each $s_{i,j}^L$ with $\alpha = 2\mu > 1$, and deduce that

$$\mathbb{E}[F(x_{k+1})] \leq F(x_0) - (1 - \frac{\mu}{2}) \frac{\gamma_{\text{low}}}{(k+2)^\mu \kappa_g^{2\mu}} \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] \\ + \left(\frac{L}{2} + 2\kappa_\Delta \right) \frac{n}{2\mu - 1} (\varsigma^{1-2\mu} - (\varsigma + \kappa_g^2(k+1))^{1-2\mu}).$$

Following the same argument as above yields that

$$\begin{aligned}
(k+1)\mathbb{E}\left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2\right] &\leq \sum_{j=0}^k \mathbb{E}[\|G_j\|^2] \\
&\leq \frac{2\kappa_g^{2\mu}(F(x_0) - F_{\text{low}})}{(1 - \frac{\mu}{2})\gamma_{\text{low}}(k+1)^{-\mu}} + \frac{n}{2\mu - 1} \left(\frac{\kappa_g^{2\mu}(L + 4\kappa_\Delta)}{\gamma_{\text{low}}(1 - \frac{\mu}{2})} \right) \times \\
&\quad \frac{\varsigma^{1-2\mu} - (\varsigma + \kappa_g^2(k+1))^{1-2\mu}}{(k+1)^{-\mu}}.
\end{aligned}$$

Rearranging the terms gives (18). \square

Note that the last fractions in the last terms of (16) and (18) have been written in a form stressing the continuity with (17), but could obviously be bounded above by the simpler

$$\frac{(\varsigma + \kappa_g^2)^{1-2\mu}}{(k+1)^\mu} \quad \text{and} \quad \frac{\varsigma^{1-2\mu}}{(k+1)^{1-\mu}}$$

respectively.

Theorem 3.5 suggests a few comments. The first is that (16), (17) and (18) guarantee the convergence of the ASGRAD algorithm with (14) to first-order critical points, because their right-hand sides all tend to zero when k tends to infinity. The rate at which this convergence occurs differs however for the three cases, depending on the parameter μ . If constants are lumped into a generic $\mathcal{O}(\cdot)$ notation, we obtain, using Jensen's inequality, that

$$\mathbb{E}\left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|\right] \leq \begin{cases} \mathcal{O}\left(\frac{1}{(k+1)^{\frac{1}{2}\mu}}\right) & (\mu \in (0, \frac{1}{2})), \\ \mathcal{O}\left(\frac{\sqrt{\log(k+1)}}{(k+1)^{\frac{1}{4}}}\right) & (\mu = \frac{1}{2}), \\ \mathcal{O}\left(\frac{1}{(k+1)^{\frac{1}{2}(1-\mu)}}\right) & (\mu \in (\frac{1}{2}, 1)). \end{cases}$$

Examining these “ k -order” bounds indicates that the best bound is that corresponding to $\mu = \frac{1}{2}$. This is nothing but the standard Adagrad algorithm.

One may then ask whether the bounds given by Theorem 3.5 are sharp. We now show that is is not the case under an additional conditional variance condition on the gradient estimator. Note that both Assumptions 3.1–3.2 automatically hold for (14) with $v_{i,j} = \varsigma + \sum_{\ell=0}^j g_{i,\ell}^2$ (for convenience, set $v_{i,-1} = \varsigma$).

Theorem 3.6. *Suppose that Assumptions 2.1–2.4 hold and that an ASGRAD algorithm is applied to problem (1) with the scaling factors defined by (14) for some $\varsigma \in (0, \kappa_g]$, $\mu \in (0, 1)$, all $i \in \{1, \dots, n\}$ and all $k \geq 0$. Suppose also that, for all $i \in \{1, \dots, n\}$ and all $k \geq 0$*

$$\text{Var}_k[g_{i,k}] = \mathbb{E}_k[g_{i,k}^2 - G_{i,k}^2] \leq \kappa_{\text{var}} G_{i,k}^2 \tag{22}$$

for some $\kappa_{\text{var}} \geq 0$. Then, for any $\theta \in (0, (1 - \frac{1}{2}\mu)\gamma_{\text{low}})$, there exists a finite $j_\theta \geq 0$ such that

$$\mathbb{E} \left[\text{average}_{j \in \{j_\theta+1, \dots, k\}} \|G_j\|^2 \right] \leq \kappa_*(\theta) \frac{(k+2)^\mu}{k-j_\theta} + \frac{n\kappa_\nabla}{k-j_\theta} \leq \frac{j_\theta+3}{(k+2)} \left(\frac{\kappa_*(\theta)}{(k+2)^{-\mu}} + n\kappa_\nabla \right), \quad (23)$$

where

$$\kappa_*(\theta) \stackrel{\text{def}}{=} \frac{\kappa_g^{2\mu}}{\theta} \{F(x_0) - F_{\text{low}} + n\kappa_\diamond\} \quad (24)$$

with

$$\begin{aligned} \kappa_\nabla &\stackrel{\text{def}}{=} \left(\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right)^{\frac{1}{\mu}}, & \kappa_\diamond &\stackrel{\text{def}}{=} \frac{\tilde{\kappa}}{\varsigma^{2\mu}} [(j_\theta+1)\kappa_g^2 + \kappa_\nabla], \\ \tilde{\kappa} &\stackrel{\text{def}}{=} \left(2\kappa_\Delta + \frac{L}{2} \right) (1 + \kappa_{\text{var}}) \left(1 + \frac{\kappa_g^2}{\varsigma} \right)^\mu, \end{aligned} \quad (25)$$

and κ_Δ defined by (11).

Proof. We have verified (at the beginning of the proof of Theorem 3.5) that the proposed scaling factors verify Assumptions 3.1 and 3.2. We invoke (2) to deduce that

$$F(x_{j+1}) \leq F(x_j) + \gamma_j G_j^\top s_j^L + \frac{L}{2} \gamma_j^2 \|s_j^L\|^2 \leq F(x_j) + \gamma_j G_j^\top s_j^L + \frac{L}{2} \|s_j^L\|^2,$$

and, taking the conditional expectation and using Lemma 3.3, that

$$\begin{aligned} \mathbb{E}_j[F(x_{j+1})] &\leq F(x_j) + \sum_{i=1}^n \mathbb{E}_j[\gamma_j G_{i,j}^\top s_{i,j}^L] + \frac{L}{2} \mathbb{E}_j[\|s_j^L\|^2], \\ &\leq F(x_j) - \sum_{i=1}^n \left(1 - \frac{\mu}{2}\right) \gamma_{\text{low}} \frac{G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + \left(2\kappa_\Delta + \frac{L}{2}\right) \mathbb{E}_j \left[\frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \right]. \end{aligned}$$

Observe now that, for each $i \in \{1, \dots, n\}$, $v_{i,j-1} + \kappa_g^2 \geq v_{i,j} \geq v_{i,j-1}$ and that $v_{i,j-1}$ is measurable knowing g_0, \dots, g_{j-1} . Hence,

$$\begin{aligned} \mathbb{E}_j[F(x_{j+1})] &\leq F(x_j) - \sum_{i=1}^n \left[1 - \frac{\mu}{2}\right] \frac{\gamma_{\text{low}} G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + \left[2\kappa_\Delta + \frac{L}{2}\right] \mathbb{E}_j \left[\frac{g_{i,j}^2}{v_{i,j-1}^{2\mu}} \right] \\ &\leq F(x_j) - \sum_{i=1}^n \left[1 - \frac{\mu}{2}\right] \frac{\gamma_{\text{low}} G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + \left[2\kappa_\Delta + \frac{L}{2}\right] \frac{\mathbb{E}_j[g_{i,j}^2]}{v_{i,j-1}^{2\mu}}. \end{aligned}$$

We now use (22) to deduce that

$$\mathbb{E}_j[F(x_{j+1})] \leq F(x_j) - \sum_{i=1}^n \left[1 - \frac{\mu}{2}\right] \frac{\gamma_{\text{low}} G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + \left[2\kappa_\Delta + \frac{L}{2}\right] (1 + \kappa_{\text{var}}) \frac{G_{i,j}^2}{v_{i,j-1}^{2\mu}}.$$

But

$$\frac{\mathbb{E}_j[v_{i,j}]}{v_{i,j-1}} \leq 1 + \frac{\kappa_g^2}{v_{i,j-1}} \leq 1 + \frac{\kappa_g^2}{\varsigma}$$

and thus

$$\frac{G_{i,j}^2}{v_{i,j-1}^{2\mu}} = \frac{(\mathbb{E}_j[v_{i,j}])^\mu}{(v_{i,j-1}\mathbb{E}_j[v_{i,j}])^\mu} \frac{G_{i,j}^2}{v_{i,j-1}^\mu} \leq \frac{(1 + \frac{\kappa_g^2}{\varsigma})^\mu}{(\mathbb{E}_j[v_{i,j}])^\mu} \frac{G_{i,j}^2}{v_{i,j-1}^\mu},$$

so that

$$\mathbb{E}_j[F(x_{j+1})] \leq F(x_j) - \sum_{i=1}^n \left[1 - \frac{\mu}{2}\right] \frac{\gamma_{\text{low}} G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} + \frac{\tilde{\kappa}}{(\mathbb{E}_j[v_{i,j}])^\mu} \frac{G_{i,j}^2}{v_{i,j-1}^\mu},$$

where $\tilde{\kappa}$ is defined by (25). Summing over iterations in $j \in \{0, \dots, k\}$ in this last inequality yields that

$$\sum_{j=0}^k \mathbb{E}_j[F(x_{j+1})] \leq \sum_{j=0}^k F(x_j) + \sum_{j=0}^k \sum_{i=1}^n \frac{G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} \left[-\gamma_{\text{low}} \left(1 - \frac{\mu}{2}\right) + \frac{\tilde{\kappa}}{v_{i,j-1}^\mu} \right]. \quad (26)$$

Now select an arbitrary $\theta \in (0, \gamma_{\text{low}}(1 - \frac{1}{2}\mu))$ and let \mathcal{I} be the (possibly empty) subset of $\{1, \dots, n\}$ such that, for all $j \geq 0$ and all $i \in \mathcal{I}$,

$$-\gamma_{\text{low}} \left(1 - \frac{\mu}{2}\right) + \frac{\tilde{\kappa}}{v_{i,j-1}^\mu} \geq -\theta.$$

This last inequality implies that, for all $j \geq 0$ and all $i \in \mathcal{I}$,

$$\mathbb{E}[v_{i,j-1}] \leq \left(\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right)^{\frac{1}{\mu}},$$

and hence, using Jensen's inequality, that, for all $k \geq 0$ and all $i \in \mathcal{I}$,

$$\begin{aligned} \mathbb{E} \left[\sum_{j=0}^k G_{i,j}^2 \right] &= \mathbb{E} \left[\sum_{j=0}^k \mathbb{E}_j[g_{i,j}]^2 \right] = \sum_{j=0}^k \mathbb{E}[g_{i,j}]^2 \\ &\leq \varsigma + \sum_{j=0}^k \mathbb{E}[g_{i,j}^2] = \mathbb{E}[v_{i,k}] \\ &\leq \left(\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right)^{\frac{1}{\mu}}. \end{aligned} \quad (27)$$

Consider now $i \in \mathcal{J} \stackrel{\text{def}}{=} \{1, \dots, n\} \setminus \mathcal{I}$ (assuming therefore that $\mathcal{J} \neq \emptyset$). For such an i , there must exist a $j_i(\theta)$ sufficiently large such that

$$-\gamma_{\text{low}} \left(1 - \frac{\mu}{2}\right) + \frac{\tilde{\kappa}}{v_{i,j-1}^\mu} \leq -\theta,$$

for $j = j_i(\theta) + 1$ and hence, since $v_{i,j}$ is an increasing function of j , for all $j \geq j_i(\theta) + 1$. If we now set

$$j_\theta = \begin{cases} \max_{i \in \mathcal{J}} j_i(\theta) & \text{if } \mathcal{J} \neq \emptyset \\ -1 & \text{otherwise.} \end{cases} \quad (28)$$

we then verify, using Assumption 2.4, that

$$\sum_{j=0}^{j_\theta} \sum_{i \in \mathcal{J}} \frac{G_{i,j}^2}{(\mathbb{E}_j[v_{i,j}])^\mu} \left(-\left(1 - \frac{\mu}{2}\right)\gamma_{\text{low}} + \tilde{\kappa} \frac{1}{v_{i,j-1}^\mu} \right) \leq n\tilde{\kappa}(j_\theta + 1) \frac{\kappa_g^2}{\zeta^{2\mu}}. \quad (29)$$

Returning to inequality (26) and using (29) for the \mathcal{J} terms and the fact that $\varsigma \leq v_{i,j} \leq (k+2)\kappa_g^2$, we deduce that

$$\begin{aligned} \sum_{j=0}^k \mathbb{E}_j[F(x_{j+1})] &\leq \sum_{j=0}^k F(x_j) - \sum_{j=j_\theta+1}^k \sum_{i \in \mathcal{J}} \frac{G_{i,j}^2}{\mathbb{E}_j[v_{i,j}]^\mu} \theta + n\tilde{\kappa}(j_\theta + 1) \frac{\kappa_g^2}{\zeta^{2\mu}} + \sum_{j=0}^k \sum_{i \in \mathcal{I}} \frac{\tilde{\kappa} G_{i,j}^2}{\zeta^{2\mu}} \\ &\leq \sum_{j=0}^k F(x_j) - \sum_{j=j_\theta+1}^k \sum_{i \in \mathcal{J}} \frac{\theta G_{i,j}^2}{\kappa_g^{2\mu} (k+2)^\mu} + \frac{\tilde{\kappa}}{\zeta^{2\mu}} \left[n(j_\theta + 1)\kappa_g^2 + \sum_{j=0}^k \sum_{i \in \mathcal{I}} G_{i,j}^2 \right]. \end{aligned}$$

We now take the full expectation and use (27) for each $i \in \mathcal{I}$ to obtain that

$$\begin{aligned} \mathbb{E}[F(x_{k+1})] &\leq F(x_0) - \frac{\theta}{\kappa_g^{2\mu} (k+2)^\mu} \mathbb{E} \left[\sum_{j=j_\theta+1}^k \sum_{i \in \mathcal{J}} G_{i,j}^2 \right] \\ &\quad + \frac{n\tilde{\kappa}}{\zeta^{2\mu}} \left[(j_\theta + 1)\kappa_g^2 + \left(\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right)^\frac{1}{\mu} \right]. \end{aligned}$$

Rearranging the terms of the last inequality and using the fact that $F(x_{k+1}) \geq F_{\text{low}}$ by Assumption 2.3 and (27) for the \mathcal{I} terms gives that

$$\begin{aligned} \mathbb{E} \left[\sum_{j=j_\theta+1}^k \|G_j\|^2 \right] &= \mathbb{E} \left[\sum_{j=j_\theta+1}^k \sum_{i \in \mathcal{J}} G_{i,j}^2 + \sum_{j=j_\theta+1}^k \sum_{i \in \mathcal{I}} G_{i,j}^2 \right] \\ &\leq \frac{\kappa_g^{2\mu} (k+2)^\mu}{\theta} \left\{ F(x_0) - F_{\text{low}} + \frac{n\tilde{\kappa}}{\zeta^{2\mu}} \left[(j_\theta + 1)\kappa_g^2 \right. \right. \\ &\quad \left. \left. + \left[\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right]^\frac{1}{\mu} \right] \right\} + n \left(\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right)^\frac{1}{\mu}. \end{aligned}$$

Dividing by $(k - j_\theta)$ gives (23)–(25). To conclude our proof, we need to examine the situation where $\mathcal{J} = \emptyset$ and (27) holds for all $i \in \mathcal{I} = \{1, \dots, n\}$. Therefore

$$(k+1) \mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq n \left(\frac{\tilde{\kappa}}{\gamma_{\text{low}}(1 - \frac{1}{2}\mu) - \theta} \right)^\frac{1}{\mu}. \quad (30)$$

We may then ignore this situation in our worst-case analysis since this last bound is clearly better than (23). \square

It is important to note that, at variance with Theorem 3.5 which states *global*² convergence rates, Theorem 3.6 only gives *asymptotic*³ rates. Indeed, our proof does not give an explicit expression of the index j_θ as a function of problem-dependent quantities only.

²That is valid for each $k \geq 0$.

³That is valid for k sufficiently large.

Our result complements that stated in Corollary 7 of [31] and Theorem 3 of [28] in that it now allows a very flexible choice of the learning rate γ_k (namely $\gamma_k \in [\gamma_{\text{low}}, 1]$) while γ_k has to be chosen as a specific function of *a priori* unknown constants⁴ in these references, if the best achievable rate of convergence is to be achieved. However, our result does not take any sparsity of the gradient sequence into account and obtains the best convergence rate at the price of the variance condition (22), which can be viewed as a “component-wise” variant of the Strong Growth Condition” (SGC) proposed in [28].

Note that expressions (24) and (23) in Theorem 3.6 have an explicit linear dependence on the problem dimension n (as in [31]), but caution should be exercised in interpreting this observation since (potentially severe) dependence on dimension may lurk in the Lipschitz constant L of Assumption 2.2.

Interestingly, Theorem 3.6 also raises the possibility (which we dismissed in the worst-case but could well positively influence the practical convergence behaviour) that $\mathcal{I} = \{1, \dots, n\}$ and (30) holds, which is then significantly better than both (16)–(18) and (23).

Even if the favourable situation we just discussed does not occur, (23) shows that (16)–(18) are not sharp whenever the conditional variance condition (22) holds. Indeed, for k sufficiently large, (23) indicates that a bound of the form

$$\mathbb{E} \left[\text{average}_{j \in \{j_\theta+1, \dots, k\}} \|G_j\| \right] = \mathcal{O} \left(\frac{1}{(k+1)^{\frac{1}{2}(1-\mu)}} \right) \quad (31)$$

is valid. Observe that this order is very close, for small μ , to that achieved by standard methods using function evaluations to enforce descent (such as steepest descent [23] or first-order trust-region [17] or adaptive regularization methods [24, 10]) in the noiseless case.

4 A “divergent series” class of ASGRAD algorithms

The key to Theorem 3.6 and its improved convergence rate is the existence of the (implicit) j_θ index. One might then wonder if another class of ASGRAD algorithms exists where such an index can be explicitly computed and a similar convergence rate achievable. This is the object of our next theorems.

Theorem 4.1. *Suppose that Assumptions 2.1–2.4 hold and that an ASGRAD algorithm is applied to problem (1) with its scaling factors being defined, for some $\nu \in (0, 1)$ and $\mu \in [\nu, \max(1, 2\nu))$, all $i \in \{1, \dots, n\}$ and all $k \geq 0$ by*

$$\rho_{i,k}(k+1)^\nu \leq w_{i,k} \leq \xi_{i,k}(k+1)^\mu, \quad (32)$$

where $\rho_{i,k}$, $w_{i,k}$ and $\xi_{i,k}$ are random variables depending on iterations $\{0, \dots, k\}$ such that $\varsigma \leq \rho_{i,k}$ and $\xi_{i,k} \leq \kappa_\xi$ for some constants $0 < \varsigma \leq \kappa_\xi$, for all $i \in \{1, \dots, n\}$ and all $k \geq 0$. Then, for $\nu \neq \frac{1}{2}$,

$$\mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq \frac{\kappa_\xi(F(x_0) - F_{\text{low}})}{\gamma_{\text{low}}(k+1)^{1-\mu}} + \frac{n L \kappa_\xi \kappa_g^2}{2(1-2\nu)\varsigma^2 \gamma_{\text{low}}} \left[\frac{1}{(k+1)^{2\nu-\mu}} - \frac{2\nu}{(k+1)^{1-\mu}} \right], \quad (33)$$

⁴Such as κ_g , L and the index of the final iteration.

while, if $\nu = \frac{1}{2}$,

$$\mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq \frac{\kappa_\xi (F(x_0) - F_{\text{low}})}{\gamma_{\text{low}} (k+1)^{1-\mu}} + \frac{n L \kappa_\xi \kappa_g^2}{2\varsigma^2 \gamma_{\text{low}}} \left(\frac{1 + \log(k+1)}{(k+1)^{1-\mu}} \right), \quad (34)$$

Proof. By using (2) and the bounds on γ_j , we derive that

$$F(x_{j+1}) \leq F(x_j) + \gamma_j G_j^T s_j^L + \frac{L}{2} \gamma_j^2 \|s_j^L\|^2 \leq F(x_j) + \gamma_j G_j^T s_j^L + \frac{L}{2} \|s_j^L\|^2. \quad (35)$$

We now derive an upper bound on the expectation $\mathbb{E}_j \left[\frac{-\gamma_j G_{i,j} g_{i,j}}{w_{i,j}} \right]$ using an argument similar to that used in the beginning of the proof of Lemma 3.3.

Consider first the case where $G_{i,j} g_{i,j} < 0$ then, $\frac{-\gamma_j G_{i,j} g_{i,j}}{w_{i,j}} \leq \frac{-G_{i,j} g_{i,j}}{\varsigma(j+1)^\nu}$ as $\gamma_j \leq 1$ and $\varsigma(j+1)^\nu \leq w_{i,j}$ (by using (32) and the bounds on $\rho_{i,k}$). Hence,

$$\mathbb{E}_j \left[\frac{-\gamma_j G_{i,j} g_{i,j}}{w_{i,j}} \right] \leq -\frac{G_{i,j}^2}{\varsigma(j+1)^\nu}. \quad (36)$$

Otherwise (i.e. if $G_{i,j} g_{i,j} \geq 0$), then, $\frac{-\gamma_j G_{i,j} g_{i,j}}{w_{i,j}} \leq \frac{-\gamma_{\text{low}} G_{i,j} g_{i,j}}{\kappa_\xi (j+1)^\mu}$ because $\gamma_{\text{low}} \leq \gamma_j$ and $w_{i,j} \leq \kappa_\xi (j+1)^\mu$ (by using (32) and the bounds on $\xi_{i,k}$). As a consequence, we have that

$$\mathbb{E}_j \left[\frac{-\gamma_j G_{i,j} g_{i,j}}{w_{i,j}} \right] \leq -\frac{\gamma_{\text{low}} G_{i,j}^2}{\kappa_\xi (j+1)^\mu}. \quad (37)$$

Defining now $\kappa_\gamma \stackrel{\text{def}}{=} \frac{\gamma_{\text{low}}}{\kappa_\xi}$, noting that $\kappa_\gamma \leq \frac{1}{\varsigma}$ and combining (36) and (37) yields that

$$\mathbb{E}_j \left[\frac{-\gamma_j G_{i,j} g_{i,j}}{w_{i,j}} \right] \leq -\frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu},$$

so that, taking the conditional expectation of (35), the last inequality and using Assumption 2.4,

$$\begin{aligned} \mathbb{E}_j[F(x_{j+1})] &\leq F(x_j) + \sum_{i=1}^n \mathbb{E}_j[\gamma_j G_{i,j} s_{i,j}^L] + \frac{L}{2} \mathbb{E}_j[\|s_j^L\|^2], \\ &\leq F(x_j) + \sum_{i=1}^n \mathbb{E}_j \left[\gamma_j G_{i,j} \frac{g_{i,j}}{w_{i,j}} \right] + \frac{L}{2} \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j}^2} \right], \\ &\leq F(x_j) - \sum_{i=1}^n \frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu} + \frac{L}{2} \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j}^2} \right], \\ &\leq F(x_j) - \sum_{i=1}^n \frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu} + \frac{L \kappa_g^2}{2} \mathbb{E}_j \left[\frac{1}{w_{i,j}^2} \right]. \end{aligned} \quad (38)$$

Using now (32) and the bounds on $\rho_{i,k}$ yields that

$$\mathbb{E}_j[F(x_{j+1})] \leq F(x_j) - \sum_{i=1}^n \kappa_\gamma \frac{G_{i,j}^2}{(j+1)^\mu} + \frac{n L \kappa_g^2}{2\varsigma^2 (j+1)^{2\nu}}.$$

Summing over all iterations from 0 to k and taking the full expectation gives that

$$\mathbb{E}[F(x_{k+1})] \leq F(x_0) - \kappa_\gamma \sum_{j=0}^k \sum_{i=1}^n \frac{\mathbb{E}[G_{i,j}^2]}{(j+1)^\mu} + \frac{nL\kappa_g^2}{2\zeta^2} \sum_{j=0}^k \frac{1}{(j+1)^{2\nu}}.$$

If we now define

$$\phi_\nu(x) \stackrel{\text{def}}{=} \begin{cases} \frac{(x+1)^{1-2\nu} - 1}{1-2\nu} & \text{if } \nu \neq \frac{1}{2} \\ \log(x+1) & \text{otherwise,} \end{cases}$$

we may bound the last inequality, using a simple sum-integral comparison and Assumption 2.3 to obtain that

$$\sum_{j=0}^k \sum_{i=1}^n \mathbb{E}[G_{i,j}^2] \leq \frac{(k+1)^\mu (F(x_0) - F_{\text{low}})}{\kappa_\gamma} + \frac{nL\kappa_g^2(k+1)^\mu}{2\zeta^2\kappa_\gamma} (1 + \phi_\nu(k)).$$

Substituting κ_γ by its value gives then,

$$\mathbb{E} \left[\text{average}_{j \in \{0, \dots, k\}} \|G_j\|^2 \right] \leq \frac{\kappa_\xi (F(x_0) - F_{\text{low}})}{\gamma_{\text{low}} (k+1)^{1-\mu}} + \frac{nL\kappa_\xi \kappa_g^2}{2\zeta^2 \gamma_{\text{low}}} \left(\frac{1 + \phi_\nu(k)}{(k+1)^{1-\mu}} \right).$$

This gives (34) when $\nu = \frac{1}{2}$. Otherwise, (33) follows by using the fact that

$$1 + \phi_\nu(k) = \frac{1}{1-2\nu} \left[\frac{1}{(k+1)^{2\nu-1}} - 2\nu \right].$$

□

The choice (32) is of course reminiscent, in a smooth but stochastic and nonconvex setting, of the “divergent stepsize” subgradient method for non-smooth convex optimization (see [1] and the many references therein), for which a $\mathcal{O}(1/\sqrt{k})$ global rate of convergence is known (Theorems 8.13 and 8.30 in this last reference).

The bounds given by Theorem 4.1 are qualitatively similar to those of Theorem 3.5, but, as in Theorem 3.6 for this case, they may be improved if we strengthen our assumptions, this time explicitly rather than implicitly.

Theorem 4.2. *Suppose that Assumptions 2.1–2.4 hold and that an ASGRAD algorithm is applied to problem (1) with its scaling factors being defined as in Theorem 4.1 with $\nu \in (0, 1)$ and $\mu \in [\nu, \max(\frac{4}{3}\nu, 1))$. Suppose additionally that the conditional variance condition (22) holds for all $i \in \{1, \dots, n\}$ and all $k \geq 0$. Then, for any $\theta \in (0, \frac{\gamma_{\text{low}}}{\kappa_\xi})$,*

$$\mathbb{E} \left[\text{average}_{j \in \{j_\theta+1, \dots, k\}} \|G_j\|^2 \right] \leq \kappa_{\#}(\theta) \frac{(k+1)^\mu}{k-j_\theta} \leq \frac{\kappa_{\#}(\theta)(j_\theta+2)}{(k+1)^{1-\mu}}, \quad (39)$$

where

$$\kappa_{\#}(\theta) \stackrel{\text{def}}{=} \frac{1}{\theta} \left(F(x_0) - F_{\text{low}} + \frac{n\kappa_g^2 L \kappa_\xi^2 (1 + \kappa_{\text{var}})}{2^{1-\mu} \zeta^4} j_\theta \right), \quad (40)$$

and

$$j_\theta \stackrel{\text{def}}{=} \left\lceil \left(\frac{L\kappa_\xi^3 (1 + \kappa_{\text{var}})}{2^{1-\mu} \zeta^4 (\gamma_{\text{low}} - \theta \kappa_\xi)} \right)^{\frac{1}{4\nu-3\mu}} \right\rceil + 1. \quad (41)$$

Proof. To simplify notation, set, for the course of this proof, $w_{i,-1} = \varsigma$, $i \in \{1, \dots, n\}$, $\frac{0}{0} = 1$ and (as in the previous theorem) $\kappa_\gamma = \gamma_{\text{low}}/\kappa_\xi$. As in the proof of Theorem 4.1, we derive (see (38)) that

$$\begin{aligned} \mathbb{E}_j[F(x_{j+1})] &\leq F(x_j) - \sum_{i=1}^n \frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu} + \frac{L}{2} \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j}^2} \right] \\ &\leq F(x_j) - \sum_{i=1}^n \frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu} + \frac{L}{2} \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j-1}^2} \left(\frac{w_{i,j-1}}{w_{i,j}} \right)^2 \right] \\ &\leq F(x_j) - \sum_{i=1}^n \frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu} + \frac{L\kappa_\xi^2 j^{2\mu}}{2\varsigma^2 j^{2\nu}} \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j-1}^2} \right] \\ &\leq F(x_j) - \sum_{i=1}^n \frac{\kappa_\gamma G_{i,j}^2}{(j+1)^\mu} + \frac{L\kappa_\xi^2 j^{2\mu}}{2\varsigma^2 j^{2\nu} w_{i,j-1}^2} (1 + \kappa_{\text{var}}) G_{i,j}^2, \end{aligned}$$

where we have used the fact that $\left(\frac{w_{i,j-1}}{w_{i,j}}\right)^2 \leq \frac{\kappa_\xi^2 j^{2\mu}}{\varsigma^2 j^{2\nu}}$ (because of (32)), the measurability of $w_{i,j-1}$ w.r.t the past and (22) to deduce the last inequality. Using now the bound $\frac{(j+1)^\mu}{w_{i,j-1}} \leq \frac{2^\mu j^\mu}{\varsigma j^\nu}$ and summing over the iterations for $j \in \{0, \dots, k\}$ yields that

$$\sum_{j=0}^k \mathbb{E}_j[F(x_{j+1})] \leq \sum_{j=0}^k F(x_j) + \sum_{j=0}^k \sum_{i=1}^n \frac{G_{i,j}^2}{(j+1)^\mu} \left(-\kappa_\gamma + \frac{\widehat{\kappa} j^{3\mu}}{w_{i,j-1} j^{3\nu}} \right), \quad (42)$$

with $\widehat{\kappa} = \frac{L\kappa_\xi^2}{2^{1-\mu}\varsigma^3} (1 + \kappa_{\text{var}})$. Note now that the definition of j_θ in (41), the fact that $4\nu > 3\mu$ and that $w_{i,j-1} \geq \varsigma j^\nu$ together imply that

$$\left(-\kappa_\gamma + \frac{\widehat{\kappa} j^{3\mu}}{w_{i,j-1} j^{3\nu}} \right) \leq -\theta, \quad (43)$$

for $j \geq j_\theta$. Hence, from (42),

$$\sum_{j=0}^k \mathbb{E}_j[F(x_{j+1})] \leq \sum_{j=0}^k F(x_j) - \theta \sum_{j=j_\theta}^k \sum_{i=1}^n \frac{G_{i,j}^2}{(j+1)^\mu} + \sum_{j=0}^{j_\theta-1} \sum_{i=1}^n \frac{G_{i,j}^2}{(j+1)^\mu} \left(-\kappa_\gamma + \frac{\widehat{\kappa} j^{3\mu}}{w_{i,j-1} j^{3\nu}} \right), \quad (44)$$

while the last term of the previous equation is bounded by

$$\sum_{j=0}^{j_\theta-1} \sum_{i=1}^n \frac{G_{i,j}^2}{(j+1)^\mu} \left(-\kappa_\gamma + \frac{\widehat{\kappa} j^{3\mu}}{w_{i,j-1} j^{3\nu}} \right) \leq \sum_{j=0}^{j_\theta-1} \sum_{i=1}^n \widehat{\kappa} \frac{G_{i,j}^2}{\varsigma} \leq \frac{n\kappa_g^2 \widehat{\kappa}}{\varsigma} j_\theta, \quad (45)$$

where we used the facts that $\|G\|_\infty \leq \kappa_g$ (because of (6)), $w_{i,j} \geq \varsigma$ (because of (32)) and $3\nu > \frac{9}{4}\mu > 2\mu$ (because of the bounds $\nu \leq \mu < \frac{4}{3}\nu$). Injecting (45) in (44), we deduce that

$$\theta \sum_{j=j_\theta}^k \sum_{i=1}^n \frac{G_{i,j}^2}{(j+1)^\mu} \leq \sum_{j=0}^k F(x_j) - \sum_{j=0}^k \mathbb{E}_j[F(x_{j+1})] + \frac{n\kappa_g^2 \widehat{\kappa}}{\varsigma} j_\theta.$$

Taking the full expectation then gives that

$$(k - j_\theta) \mathbb{E} \left[\text{average}_{j \in \{j_\theta+1, \dots, k\}} \|G_j\|^2 \right] \leq \mathbb{E} \left[\sum_{j=j_\theta}^k \sum_{i=1}^n G_{i,j}^2 \right] \leq \frac{(k+1)^\mu}{\theta} \left[F(x_0) - F_{\text{low}} + \frac{n\kappa_g^2 \widehat{\kappa}}{\varsigma} j_\theta \right]. \quad (46)$$

which gives the desired result. \square

The k -order of convergence of $\mathbb{E} \left[\text{average}_{j \in \{j_\theta+1, \dots, k\}} \|G_j\| \right]$ implied by (39) is therefore

$$\mathcal{O} \left(\frac{1}{(k+1)^{\frac{1}{2}(1-\mu)}} \right)$$

which is identical to (31), except that now (41) now gives an explicit formula for j_θ , thereby quantifying what is meant by the “ k sufficiently large” assumption used to derive (31).

It is of course possible to combine the ideas of the two ASGRAD classes considered so far, for instance by defining both $\xi_{i,k}$ and $\rho_{i,k}$ in (32) to be equal to $\varsigma + \sum_{j=0}^k g_{i,j}^2 / (k+1)$, without altering the results of Theorem 4.2.

5 Numerical experiments on deep-learning datasets

We now provide some numerical illustration of the algorithmic variants discussed in the previous sections, in the context of deep-learning using neural networks for image processing. The problem considered is to classify images in a preset number of classes. A neural network is then constructed whose output is fed in a cross-entropy objective function⁵ (also called “loss” in the deep-learning jargon). This a relatively standard setting in which noise arises from sampling the terms of the finite sum defining the objective function.

We trained two network architectures, namely the simple `cifar-nv` convolutional network of [16] and a small `resnet18` model [18], on four standard datasets of 32×32 images: CIFAR10 and CIFAR100⁶, SVHN⁷ and FMNIST [30]. For these experiments, we used haiku [19] and optax [20] two JAX [9] based libraries on a workstation with four GTX 1080TI. We now compare the numerical performance of (14) for various μ values in (0.1, 0.5, 0.9) and of two scaling factor choices satisfying (32). In order to define these, first set

$$\xi_k = \max(\varsigma, \xi_{k-1}, |g_k|), \quad (k \geq 0), \quad (47)$$

with $\xi_{-1} = 0$ and the max being understood componentwise. They are then defined by

$$w_{i,k} = \xi_k (k+1)^\nu \quad (48)$$

and

$$w_{i,k} = \max\left(\varsigma, \frac{1}{k+1} \sum_{j=0}^k |g_{i,j}|\right) (k+1)^\nu. \quad (49)$$

These two scalings, denoted respectively by *maxgi* and *avrgi*, verify (32) with $\varsigma = \varsigma$ and $\kappa_\xi = \kappa_g$. In (48) and (49), we chose $\nu = 0.1$ and $\varsigma = 0.01$. The corresponding variants are

⁵<https://pytorch.org/docs/stable/generated/torch.nn.CrossEntropyLoss.html>

⁶<https://www.cs.toronto.edu/~kriz/cifar.html>

⁷<http://ufldl.stanford.edu/housenumbers>

called *maxgi* and *avrgi*, respectively. Note that the scaling factors are increasing in *maxgi* but no necessarily so in *avrgi*.

For all experiments, we also chose a fixed⁸ learning rate policy with $\gamma_k = \gamma = 5 \cdot \{10^{-4}, 10^{-5}\}$ for all $k \geq 0$. We used the same random initialization (as specified by haiku) for all scaling choices and followed the data-augmentation procedure of [16], both for training and testing. We trained the models for a total of 100000 steps with a batchsize (sample size) of 128 using the cross-entropy loss function. We report the training and test accuracies (the latter on a sample of size 128 from the test dataset) every 500 steps.

The results of these experiments (averaged over three random runs) are presented in Figures 1–4 for the CIFAR10 and CIFAR100 datasets. The corresponding results for SVHN and FMNIST are reported in Appendix. In each figure, the top panels show the evolution of the training accuracy (as a function of the number of steps), and the bottom panels that of the test accuracy. The average values are shown as thick lines and the shaded areas of corresponding colour give the 67% confidence intervals.




Figure 1: CIFAR10: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5 \cdot 10^{-4}$ (left) and $\gamma = 5 \cdot 10^{-5}$ (right) on the *cifar-nv* architecture

These experiments are obviously not meant to replace significant numerical testing, but, albeit caution must be exercised not to extrapolate from limited data, they still suggest a few tentative comments.

- For fixed learning rates, the methods *maxgi* and *avrgi* of the second ASGRAD class (introduced in Section 4) seem to produce relatively good results on our example, both in training and testing, often outperforming the Adagrad-like variants of the first class (of Section 3), sometimes more than marginally.
- The relative behaviour of the tested variants does not differ significantly between the two tested network architectures, even if the test accuracy is (as expected) slightly lower for the *resnet18* case.

⁸Our choice of a fixed learning rate policy is meant to focus on the intrinsic properties of each scaling factor option.




Figure 2: CIFAR10: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5.10^{-4}$ (left) and $\gamma = 5.10^{-5}$ (right) on the **resnet18** architecture




Figure 3: CIFAR100: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5.10^{-4}$ (left) and $\gamma = 5.10^{-5}$ (right) on the **cifar-nv** architecture



Figure 4: CIFAR100: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5 \cdot 10^{-4}$ (left) and $\gamma = 5 \cdot 10^{-5}$ (right) on the *resnet18* architecture

- Among Adagrad-like variants of the first class, those with a larger μ handle smaller learning rates better on these examples, a behaviour admittedly not predicted by our theory.
- The comparison of left and right panels in each figure unsurprisingly shows that, albeit our theory does not depend on the choice of $\gamma_k \in (0, 1]$, the practical convergence behaviour may be affected by this choice (and other factors such as finite-sum sampling strategy).

These conclusions are supported and reinforced by the results for SVHN and FMNIST in appendix.

6 Conclusions

We have introduced a first-order trust-region framework for minimization methods and derived complexity upper bounds for two classes of interest, the first containing the standard Adagrad. These bounds give the best complexity to Adagrad in the first class. We have also shown these bounds can be improved for both classes under an additional variance condition, in which case the parameter choice yielding the best bounds no longer corresponds to Adagrad. This improvement is asymptotic and implicit for the first class and explicit for the second.

Our numerical illustrations of the discussed methods on examples arising from deep-learning applications indicate that methods of the second class have merits, but also that, at least in our examples, there remains some distance from the above theory to real behaviour. This may possibly be because the complexity bounds may not be sharp, but also, fortunately, because the worst-case happens very rarely in practice.

Acknowledgments This work was supported in part by 3IA Artificial and Natural Intelligence Toulouse Institute, French "Investing for the Future - PIA3" program under the Grant agreement ANR-19-PI3A-0004". The experiments presented in this paper were carried out on the OSIRIM platform administered by IRIT and supported by CNRS, the Region Midi-Pyrénées, the French Government, and ERDF (see <http://osirim.irit.fr/site/en>).

References

- [1] A. Beck. *First-order Methods in Optimization*. Number 25 in MOS-SIAM Optimization Series. SIAM, Philadelphia, USA, 2017.
- [2] S. Bellavia, G. Gurioli, B. Morini, and Ph. L. Toint. Adaptive regularization algorithms with inexact evaluations for nonconvex optimization. *SIAM Journal on Optimization*, 29(4):2881–2915, 2019.
- [3] S. Bellavia, G. Gurioli, B. Morini, and Ph. L. Toint. A stochastic ARC method with inexact function and random derivatives evaluations. In *Proceedings of the International Conference on Machine Learning (ICML2020)*, 2020.
- [4] S. Bellavia, G. Gurioli, B. Morini, and Ph. L. Toint. The impact of noise on evaluation complexity: The deterministic trust-region case. arXiv:2104.02519, 2021.
- [5] S. Bellavia, G. Gurioli, B. Morini, and Ph. L. Toint. Quadratic and cubic regularization methods with inexact function and random derivatives for finite-sum minimization. In *Proceedings of the ICCSA 2021*, 2021.
- [6] S. Bellavia, G. Gurioli, B. Morini, and Ph. L. Toint. Adaptive regularization algorithm for nonconvex optimization using inexact function evaluations and randomly perturbed derivatives. *Journal of Complexity*, 68, 2022.
- [7] A. Berahas, L. Cao, and K. Scheinberg. Global convergence rate analysis of a generic line search algorithm with noise. *SIAM Journal on Optimization*, 31:1489–1518, 2021.
- [8] J. Blanchet, C. Cartis, M. Menickelly, and K. Scheinberg. Convergence rate analysis of a stochastic trust region method via supermartingales. *INFORMS Journal on Optimization*, 1(2):92–119, 2019.
- [9] J. Bradbury, R. Frostig, P. Hawkins, M.-J. Johnson, C. Leary, D. Maclaurin, G. Necula, A. Paszke, J. VanderPlas, S. Wanderman-Milne, and Q. Zhang. JAX: composable transformations of Python+NumPy programs, 2018.
- [10] C. Cartis, N. I. M. Gould, and Ph. L. Toint. Adaptive cubic overestimation methods for unconstrained optimization. Part II: worst-case function-evaluation complexity. *Mathematical Programming, Series A*, 130(2):295–319, 2011.
- [11] C. Cartis, N. I. M. Gould, and Ph. L. Toint. *Evaluation complexity of algorithms for nonconvex optimization*. Number 30 in MOS-SIAM Series on Optimization. SIAM, Philadelphia, USA, June 2022.
- [12] C. Cartis and K. Scheinberg. Global convergence rate analysis of unconstrained optimization methods based on probabilistic models. *Mathematical Programming, Series A*, 159(2):337–375, 2018.
- [13] R. Chen, M. Menickelly, and K. Scheinberg. Stochastic optimization using a trust-region method and random models. *Mathematical Programming, Series A*, 169(2):447–487, 2018.
- [14] A. Défossez, L. Bottou, F. Bach, and N. Usunier. A simple convergence proof for Adam and Adagrad. arXiv:2003.02395v2, 2020.
- [15] J. Duchi, E. Hazan, and Y. Singer. Adaptive subgradient methods for online learning and stochastic optimization. *Journal of Machine Learning Research*, 12, July 2011.
- [16] I. Gitman and B. Ginsburg. Comparison of batch normalization and weight normalization algorithms for the large-scale image classification. arXiv:1709.08145v1, 2017.
- [17] S. Gratton, A. Sartenaer, and Ph. L. Toint. Recursive trust-region methods for multiscale nonlinear optimization. *SIAM Journal on Optimization*, 19(1):414–444, 2008.
- [18] K. He, X. Zhang, S. Ren, and J. Sun. Deep residual learning for image recognition. arXiv:1512.03385, 2015.
- [19] T. Hennigan, T. Cai, T. Norman, and I. Babuschkin. Haiku: Sonnet for JAX, 2020.

- [20] M. Hessel, D. Budden, F. Viola, M. Rosca, E. Sezener, and T. Hennigan. Optax: composable gradient transformation and optimisation, in JAX!, 2020.
- [21] D. Kingma and J. Ba. Adam: A method for stochastic optimization. In *Proceedings in the International Conference on Learning Representations (ICLR)*, 2015.
- [22] X. Li and F. Orabona. On the convergence of stochastic gradient descent with adaptive stepsizes. *AI Stats*, 2019.
- [23] Yu. Nesterov. *Introductory Lectures on Convex Optimization*. Applied Optimization. Kluwer Academic Publishers, Dordrecht, The Netherlands, 2004.
- [24] Yu. Nesterov and B. T. Polyak. Cubic regularization of Newton method and its global performance. *Mathematical Programming, Series A*, 108(1):177–205, 2006.
- [25] C. Paquette and K. Scheinberg. A stochastic line search method with convergence rate analysis. *SIAM Journal on Optimization*, 30(1):349–376, 2020.
- [26] S. Reddi, S. Kale, and S. Kumar. On the convergence of Adam and beyond. In *Proceedings in the International Conference on Learning Representations (ICLR)*, 2018.
- [27] T. Tieleman and G. Hinton. Lecture 6.5-RMSPROP. COURSERA: Neural Networks for Machine Learning, 2012.
- [28] S. Vaswani, F. Bach, and M. Schmidt. Fast and faster convergence of SGD for over-parameterized models (and an accelerated perceptron). In *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics (AISTATS) 2019, Naha, Okinawa, Japan*, volume 89, 2019.
- [29] R. Ward, X. Wu, and L. Bottou. Adagrad stepsizes: sharp convergence over nonconvex landscapes. In *Proceedings in the International Conference on Machine Learning (ICML2019)*, 2019.
- [30] H. Xiao, K. Rasul, and R. Vollgraf. Fashion-mnist: a novel image dataset for benchmarking machine learning algorithms. arXiv:1708.07747, 2017.
- [31] D. Zhou, Y. Tang, Z. Yang, Y. Cao, and Q. Gu. On the convergence of adaptive gradient methods for nonconvex optimization. In *Proceedings of OPT2020: 12th Annual Workshop on Optimization for Machine Learning*, 2020.
- [32] F. Zou, L. Shen, Z. Jie, J. Sun, and W Liu. A sufficient condition for convergences of Adam and RMSprop. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, 2019.

A first technical lemma

Lemma .1. *Let $\mu \in (0, 1]$. Let $x, y \in \mathbb{R}^+ \setminus \{0\}$. Then*

$$\frac{|x^\mu - y^\mu|}{x^\mu y^\mu} \leq \mu \frac{|x - y|}{xy^\mu} + \mu \frac{|x - y|}{x^\mu y}. \quad (50)$$

Proof. Let us first consider the case $x \geq y$. Remembering that $u^\mu \leq 1 + \mu(u - 1)$ for $u > 0$ and taking $u = \frac{x}{y}$, we successively derive that

$$\begin{aligned} \frac{x^\mu}{y^\mu} &\leq 1 + \mu \left(\frac{x}{y} - 1 \right), \\ x^\mu - y^\mu &\leq \mu \left(\frac{xy^\mu}{y} - y^\mu \right), \\ x^\mu - y^\mu &\leq \mu y^{\mu-1} (x - y), \\ \frac{x^\mu - y^\mu}{x^\mu y^\mu} &\leq \mu \frac{x - y}{x^\mu y}. \end{aligned} \quad (51)$$

Hence the inequality (50) is valid when $x \geq y$. For the symmetric case ($y \geq x$), we similarly obtain that

$$\frac{y^\mu - x^\mu}{x^\mu y^\mu} \leq \mu \frac{y - x}{y^\mu x}. \quad (52)$$

Combining (51) and (52) yields the desired result.

Proof of Lemma 3.3

Let us consider an iteration index $j \geq 0$ and a component index $i \in \{1, \dots, n\}$. We first use the definition of s^L in (4) and the fact that $w_{i,j} = v_{i,j}^\mu$ (Assumption 3.1) to obtain that

$$\mathbb{E}_j [\gamma_j G_{i,j} s_{i,j}^L] = -\mathbb{E}_j \left[\gamma_j \frac{G_{i,j} g_{i,j}}{v_{i,j}^\mu} \right] = -\mathbb{E}_j \left[\gamma_j \frac{G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \right] + \mathbb{E}_j \left[\gamma_j G_{i,j} g_{i,j} \left(\frac{1}{\mathbb{E}_j[v_{i,j}^\mu]} - \frac{1}{v_{i,j}^\mu} \right) \right]. \quad (53)$$

If $G_{i,j} g_{i,j} \geq 0$, then

$$-\frac{\gamma_j G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \leq -\gamma_{\text{low}} \frac{G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]}$$

and thus

$$\mathbb{E}_j \left[-\frac{\gamma_j G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \right] \leq -\gamma_{\text{low}} \frac{G_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \mathbb{E}_j[g_{i,j}] = -\gamma_{\text{low}} \frac{G_{i,j}^2}{\mathbb{E}_j[v_{i,j}^\mu]}. \quad (54)$$

Otherwise, if $G_{i,j} g_{i,j} < 0$, we may deduce that

$$-\frac{\gamma_j G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \leq -\frac{G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]},$$

implying that

$$\mathbb{E}_j \left[-\frac{\gamma_j G_{i,j} g_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \right] \leq -\frac{G_{i,j}}{\mathbb{E}_j[v_{i,j}^\mu]} \mathbb{E}_j[g_{i,j}] = -\frac{G_{i,j}^2}{\mathbb{E}_j[v_{i,j}^\mu]} \leq -\gamma_{\text{low}} \frac{G_{i,j}^2}{\mathbb{E}_j[v_{i,j}^\mu]}. \quad (55)$$

Combining (53), (54) and (55) gives that

$$\mathbb{E}_j[\gamma_j G_{i,j} s_{i,j}^L] \leq -\gamma_{\text{low}} \frac{G_{i,j}^2}{\mathbb{E}_j[v_{i,j}]^\mu} + \mathbb{E}_j \left[\underbrace{\gamma_j G_{i,j} g_{i,j} \frac{v_{i,j}^\mu - \mathbb{E}_j[v_{i,j}]^\mu}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]^\mu}}_A \right]. \quad (56)$$

We now derive an upper bound on the absolute value of the A term by successively using Lemma (.1), Assumption 3.1 and the bound $\gamma_j \leq 1$ to obtain that

$$\begin{aligned} |A| &= |\gamma_j G_{i,j} g_{i,j}| \frac{|v_{i,j}^\mu - \mathbb{E}_j[v_{i,j}]^\mu|}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]^\mu} \\ &\leq \mu |\gamma_j G_{i,j} g_{i,j}| \frac{|v_{i,j} - \mathbb{E}_j[v_{i,j}]|}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]} + \mu |\gamma_j G_{i,j} g_{i,j}| \frac{|v_{i,j} - \mathbb{E}_j[v_{i,j}]|}{v_{i,j} \mathbb{E}_j[v_{i,j}]^\mu} \\ &\leq \underbrace{\mu |G_{i,j} g_{i,j}| \kappa_v \frac{\mathbb{E}_j[g_{i,j}^2]}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]}}_B + \underbrace{\mu |G_{i,j} g_{i,j}| \kappa_v \frac{g_{i,j}^2}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]}}_C \\ &\quad + \underbrace{\mu |G_{i,j} g_{i,j}| \kappa_v \frac{\mathbb{E}_j[g_{i,j}^2]}{v_{i,j} \mathbb{E}_j[v_{i,j}]^\mu}}_D + \underbrace{\mu |G_{i,j} g_{i,j}| \kappa_v \frac{g_{i,j}^2}{v_{i,j} \mathbb{E}_j[v_{i,j}]^\mu}}_E. \end{aligned}$$

We now use Young's inequality with $p = q = 2$, that is

$$\forall \lambda > 0, x, y \in \mathbb{R}^+, xy \leq \frac{\lambda}{2} x^2 + \frac{y^2}{2\lambda}, \quad (57)$$

to successively handle the 4 terms of the last bound.

- For the first term B , we choose

$$x = \frac{|G_{i,j}|}{\mathbb{E}_j[v_{i,j}]^\mu}, \quad \lambda = \frac{\gamma_{\text{low}} \mathbb{E}_j[v_{i,j}]^\mu}{4} \quad \text{and} \quad y = \kappa_v |g_{i,j}| \frac{\mathbb{E}_j[g_{i,j}^2]}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]^{1-\mu}}.$$

Using (57), Assumptions 2.4, 3.2 and (9), we obtain that

$$\begin{aligned} B &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \frac{\mathbb{E}_j[g_{i,j}^2]^2}{\mathbb{E}_j[v_{i,j}]^{2-\mu}}, \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \mathbb{E}_j[g_{i,j}^2]^\mu \frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \kappa_g^{2\mu} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}}. \end{aligned}$$

Taking now the expectation over $\mathbb{E}_j[\cdot]$ yields that

$$\mathbb{E}_j[B] \leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \kappa_g^{2\mu} \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j}^2} \right]. \quad (58)$$

- Now consider the C term. In this case, we choose

$$x = \frac{|G_{i,j}g_{i,j}|}{\mathbb{E}_j[v_{i,j}]^\mu}, \quad \lambda = \gamma_{\text{low}} \frac{\mathbb{E}_j[v_{i,j}]^\mu}{4\mathbb{E}_j[g_{i,j}^2]} \quad \text{and} \quad y = \kappa_v \frac{g_{i,j}^2}{v_{i,j}^\mu \mathbb{E}_j[v_{i,j}]^{1-\mu}}$$

to deduce from (57) that

$$\begin{aligned} C &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} \frac{g_{i,j}^2}{\mathbb{E}_j[g_{i,j}^2]} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{g_{i,j}^4}{v_{i,j}^{2\mu}} \frac{\mathbb{E}_j[g_{i,j}^2]}{\mathbb{E}_j[v_{i,j}]^{2-\mu}} \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} \frac{g_{i,j}^2}{\mathbb{E}_j[g_{i,j}^2]} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{\kappa_g^2 g_{i,j}^2}{v_{i,j}^{2\mu}} \frac{1}{\mathbb{E}_j[v_{i,j}]^{1-\mu}} \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} \frac{g_{i,j}^2}{\mathbb{E}_j[g_{i,j}^2]} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{\kappa_g^2}{\varsigma_{\min}^{1-\mu}} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}}, \end{aligned}$$

where we successively used the facts that $\mathbb{E}_j[g_{i,j}^2] \leq \mathbb{E}_j[v_{i,j}]$ (because of (9)), $g_{i,j}^2 \leq \kappa_g^2$ (because of Assumption 2.4) and $\mathbb{E}_j[v_{i,j}]^{1-\mu} \geq \varsigma_{\min}^{1-\mu}$ (because of (8)). Taking the expectation over $\mathbb{E}_j[\cdot]$ then gives that

$$\mathbb{E}_j[C] \leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{\kappa_g^2}{\varsigma_{\min}^{1-\mu}} \mathbb{E}_j \left[\frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \right]. \quad (59)$$

(Note that we can divide by $\mathbb{E}_j[g_{i,j}^2]$ above, as it suffice to notice that $\mathbb{E}_j[g_{i,j}^2] = 0$ implies $g_{i,j}^2 = 0$. C would then be equal to zero and (59) would still be verified.)

- Let us now handle the D term. Choosing

$$x = \frac{|G_{i,j}|}{\mathbb{E}_j[v_{i,j}]^\mu}, \quad \lambda = \gamma_{\text{low}} \frac{\mathbb{E}_j[v_{i,j}]^\mu}{4} \quad \text{and} \quad y = \kappa_v |g_{i,j}| \frac{\mathbb{E}_j[g_{i,j}^2]}{v_{i,j}},$$

we now deduce from (57) that

$$\begin{aligned} D &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{g_{i,j}^2 \mathbb{E}_j[g_{i,j}^2]^2}{\mathbb{E}_j[v_{i,j}]^\mu v_{i,j}^2}, \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \frac{1}{v_{i,j}^{2-2\mu}} \frac{\mathbb{E}_j[g_{i,j}^2]^2}{\mathbb{E}_j[v_{i,j}]^\mu} \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \frac{1}{v_{i,j}^{2-2\mu}} \mathbb{E}_j[g_{i,j}^2]^{2-\mu} \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{\kappa_g^{4-2\mu}}{\varsigma_{\min}^{2-2\mu}} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}}, \end{aligned}$$

where, as for the C term, we used the facts that $\mathbb{E}_j[g_{i,j}^2]^\mu \leq \mathbb{E}_j[v_{i,j}]^\mu$, $g_{i,j}^2 \leq \kappa_g^2$ and $v_{i,j}^{2-2\mu} \geq \varsigma_{\min}^{2-2\mu}$. Taking the expectation $\mathbb{E}_j[\cdot]$ yields, in this case, that

$$\mathbb{E}_j[D] \leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{\kappa_g^{4-2\mu}}{\varsigma_{\min}^{2-2\mu}} \mathbb{E}_j \left[\frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \right]. \quad (60)$$

- Finally consider the E term. Choosing

$$x = \frac{|G_{i,j}g_{i,j}|}{\mathbb{E}_j[v_{i,j}]^\mu}, \quad \lambda = \gamma_{\text{low}} \frac{\mathbb{E}_j[v_{i,j}]^\mu}{4\mathbb{E}_j[g_{i,j}^2]} \quad \text{and} \quad y = \kappa_v \frac{g_{i,j}^2}{v_{i,j}}$$

in (57) then gives that

$$\begin{aligned} \mathbb{E}_j[E] &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} \frac{g_{i,j}^2}{\mathbb{E}_j[g_{i,j}^2]} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \frac{g_{i,j}^4 \mathbb{E}_j[g_{i,j}^2]}{\mathbb{E}_j[v_{i,j}]^\mu v_{i,j}^2} \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} \frac{g_{i,j}^2}{\mathbb{E}_j[g_{i,j}^2]} \\ &\quad + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \mathbb{E}_j[g_{i,j}^2]^{1-\mu} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \left(\frac{1}{v_{i,j}^{1-2\mu}} \mathbf{1}_{\mu < \frac{1}{2}} + \frac{|g_{i,j}^{4-4\mu}|}{v_{i,j}^{2-2\mu}} |g_{i,j}^{4\mu-2}| \mathbf{1}_{\mu \geq \frac{1}{2}} \right) \\ &\leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} \frac{g_{i,j}^2}{\mathbb{E}_j[g_{i,j}^2]} + 2 \frac{\kappa_v^2}{\gamma_{\text{low}}} \kappa_g^{2-2\mu} \frac{g_{i,j}^2}{v_{i,j}^{2\mu}} \left(\frac{1}{\varsigma_{\min}^{1-2\mu}} \mathbf{1}_{\mu < \frac{1}{2}} + \kappa_g^{4\mu-2} \mathbf{1}_{\mu \geq \frac{1}{2}} \right), \end{aligned}$$

where we once more used the facts that $\mathbb{E}_j[g_{i,j}^2]^\mu \leq \mathbb{E}_j[v_{i,j}]^\mu$ and $|g_{i,j}| \leq \kappa_g$, in turn implying that

$$g_{i,j}^2 \leq v_{i,j} \quad \text{and} \quad v_{i,j} \geq \varsigma_{\min} \quad \text{if} \quad \mu < \frac{1}{2}$$

and

$$|g_{i,j}^{4-4\mu}| \leq v_{i,j}^{2-2\mu} \quad \text{and} \quad |g_{i,j}^{4\mu-2}| \leq \kappa_g^{4\mu-2} \quad \text{if} \quad \mu \geq \frac{1}{2}.$$

Taking the expectation $\mathbb{E}_j[\cdot]$, we deduce that

$$\mathbb{E}_j[E] \leq \gamma_{\text{low}} \frac{G_{i,j}^2}{8\mathbb{E}_j[v_{i,j}]^\mu} + \frac{\kappa_v^2}{\gamma_{\text{low}}} \kappa_g^{2-2\mu} \left(\frac{1}{\varsigma_{\min}^{1-2\mu}} \mathbf{1}_{\mu < \frac{1}{2}} + \kappa_g^{4\mu-2} \mathbf{1}_{\mu \geq \frac{1}{2}} \right) \mathbb{E}_j \left[\frac{g_{i,j}^2}{w_{i,j}^2} \right]. \quad (61)$$

Summing now (58), (59), (60) and (61) and substituting the obtained upper-bound of A in (56), we finally obtain (10) with (11).

Proof of Lemma 3.4

Consider first the case where $\alpha \neq 1$ and note that $\frac{1}{(1-\alpha)}x^{1-\alpha}$ is then a non-decreasing and concave function on $(0, +\infty)$. Setting $b_{-1} = 0$ and using these properties, we obtain that, for $j \geq 0$,

$$\begin{aligned} \frac{a_j}{(\varsigma + b_j)^\alpha} &\leq \frac{1}{1-\alpha} \left((\varsigma + b_j)^{1-\alpha} - (\varsigma + b_j - a_j)^{1-\alpha} \right) \\ &\leq \frac{1}{1-\alpha} \left((\varsigma + b_j)^{1-\alpha} - (\varsigma + b_{j-1})^{1-\alpha} \right). \end{aligned}$$

We then obtain (12) by summing this inequality for $j \in \{0, \dots, k\}$.

Suppose now that $\alpha = 1$, We then use the concavity and non-decreasing character of the logarithm to derive that

$$\frac{a_j}{(\zeta + b_j)^\alpha} = \frac{a_j}{\zeta + b_j} \leq \log(\zeta + b_j) - \log(\zeta + b_j - a_j) \leq \log(\zeta + b_j) - \log(\zeta + b_{j-1}).$$

The inequality (13) then again follows by summing for $j \in \{0, \dots, k\}$.

Results for SVHN and FMNIST




Figure 5: SVHN: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5.10^{-4}$ (left) and $\gamma = 5.10^{-5}$ (right) on the *cifar-nv* architecture




Figure 6: SVHN: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5.10^{-4}$ (left) and $\gamma = 5.10^{-5}$ (right) on the *resnet18* architecture




Figure 7: FMNIST: Training (top) and test (bottom) accuracies for the Adagrad-like ($\mu \in (0.1, 0.5, 0.9)$), *maxgi* and *avrgi* variants with $\gamma = 5.10^{-4}$ (left) and $\gamma = 5.10^{-5}$ (right) on the **resnet18** architecture